



Ministry of Tourism & Wildlife



NATIONAL WILDLIFE CENSUS 2021 REPORT

Abridged Version



Wildlife
Research
& Training
Institute



ISBN 978-9914-40-397-8



Written and Edited By (Authors): Brig. (Rtd) John Waweru², Dr. Patrick Omondi¹, Dr. Shadrack Ngene¹, Dr. Joseph Mukeka¹, Edwin Wanyonyi², Bernard Ngoru¹, Stephen Mwiu¹, Daniel Muteti¹, Fredrick Lala¹, Linus Kariuki¹, Dr. Festus Ihwagi³, Sospeter Kiambi¹, Cedric Khyale¹, Geoffrey Bundotich¹, Dr. Fred Omengo¹, Peter Hongo¹, Peter Maina¹, Faith Muchiri¹, Dr. Mohammed Omar¹, Judith Nyunja¹, Joseph Edebe¹, James Mathenge¹, George Anyona¹, Chrispin Ngesa¹, John Gathua⁴, Lucy Njino⁴, Gakami Njenga⁴, Anthony Wandera⁵, Samuel Mutisya⁶, Redempta Njeri⁷, David Kimanzi⁸, Titus Imboma⁹, Jane Wambugu¹, Timoth Mwinami⁹, Ali Kaka¹⁰ and Dr. Erustus Kanga¹⁰

¹Wildlife Research and Training Institute, P.O. Box 842-20117, Naivasha, Kenya

²Kenya Wildlife Service, P.O. Box 409241-00100, Nairobi, Kenya

³Save the elephants, P.O. Box 54667-00200, Nairobi, Kenya

⁴Directorate of Resource Surveys and Remote Sensing, P.O. Box 47146-00100, Nairobi, Kenya

⁵Northern Rangeland Trust, Private Bag-60300, Isiolo, Kenya

⁶OI Pajeta Conservancy, Private Bag-10400, Nanyuki, Kenya

⁷Space for Giants, P.O. Box 174-10400, Nanyuki, Kenya

⁸Mara Elephant Project, P.O. Box 2606-00502, Narok, Kenya

⁹National Museums of Kenya, P.O. Box 40658-00100, Nairobi, Kenya

¹⁰Ministry of Tourism and Wildlife, P.O. Box 30126-00100, Nairobi, Kenya

Concept and Design by: Boniface Gor, www.digimatt.co.ke @digimattsol

©**Photos:** David Macharia/Versatile Photography | David Gotlib | KWS | Shutterstock | Freepik

Published in July 2021 by the Wildlife Research and Training Institute (WRTI) and Kenya Wildlife Service (KWS)

Any Reproduction in Full or in Part Must Mention the Title and Credit the Above-Mentioned Publisher as The Copyright Owners.

©Text July 2021 WRTI & KWS

All Rights Reserved



CONT


LIST OF FIGURES	v
ABBREVIATIONS	vi
EXECUTIVE SUMMARY	vii
MESSAGE FROM THE PRESIDENT	ix
FOREWORD	x
PREFACE	xi
ACKNOWLEDGEMENTS	xii

1.0 GENERAL INTRODUCTION

1.1 Background	2
1.2 Goal and Objectives	3
1.3 Project Justification	3

2.0 MATERIALS & METHODS

2.1 Study Areas	6
2.2 Methodology	7



3.0 THE 2020 WILDLIFE CENSUS RESULTS SUMMARY

WILDLIFE NUMBERS AT A GLANCE 10

3.1	KEY WILDLIFE SPECIES	15
3.1.1	Elephant	17
3.1.2	Rhino	21
3.1.3	Buffalo	24
3.1.4	Giraffe	27
3.1.5	Grevy's zebra	30
3.1.6	Sable Antelope	33
3.1.7	Mountain Bongo	36
3.1.8	Roan Antelope	39
3.1.9	Sitatunga Antelope	42
3.1.10	Hirola Antelope	45
3.1.11	The African Lion	48
3.1.12	Spotted Hyena	51
3.1.13	Cheetah	54
3.1.14	Wild Dog	57
3.1.15	Waterfowls	60



CONTENTS

3.0 THE 2020 WILDLIFE CENSUS RESULTS SUMMARY

3.1 RESULTS BY ECOSYSTEMS

3.2.1	Tsavo Ecosystem	64
3.2.2	Laikipia-Samburu-Marsabit-Meru Ecosystem	66
3.2.3	Maasai Mara National Reserve (MMNR) - Ecosystem	69
3.2.4	Amboseli-Magadi ecosystem	71
3.2.5	Athi -Kapiti Ecosystem	73
3.2.6	Lamu-Lower Garissa Ecosystem	75
3.2.7	Naivasha-Nakuru Ranches	77
3.2.8	East Isiolo County	79
3.2.9	Lake Nakuru National Park	81
3.2.10	Nairobi National Park	83
3.2.11	Oldonyo Sabuk National Park	85
3.2.12	Mwea National Reserve	87
3.2.13	Ruma National Park	89
3.2.14	Shimba Hills Conservation Areas	91
3.2.15	Tana River Remnant Forests	93
3.2.16	Mandera County Ecosystem	95
3.2.17	Garissa County	97
3.2.18	Wajir County	99
3.2.19	Turkana County	101
3.2.20	Marine Ecosystem	103
3.2.21	Nasalot-South Turkana-Kerio Valley ecosystem	105

4.0 CONCLUSIONS & RECOMMENDATIONS

4.1	CONCLUSIONS	110
4.2	RECOMMENDATIONS	111
	REFERENCES	112





LIST OF FIGURES

Figure 1:	A Map Showing the Kenya Wildlife Service Conservation Area Where the Census took Place between April and July 2021	6
Figure 2:	The Kenya National Elephant Range as at July 2021	19
Figure 3:	Map Showing Rhino Conservation Areas	23
Figure 4:	Buffalo Population Trends in Kenya	26
Figure 5:	Distribution of Giraffes in Kenya	29
Figure 6:	Grevy'S Zebra Distribution Map in Kenya	32
Figure 7:	Historical Distribution of Roosevelt's Sable Antelope in Kenya from 1884 to Present (Source: Modified from Butynski et al., 2015)	35
Figure 8:	Map of Current Mountain Bongo Range in Kenya	38
Figure 9:	Roan Distribution in Ruma N.P	41
Figure 10:	Roan Antelope Sighting in Ruma NP	41
Figure 11:	Distribution of Sitatunga's in Western conservation Area	44
Figure 12:	Distribution of Sitatungas in Central Rift Conservation Area	44
Figure 13:	Map Showing the Distribution of Hirola in Kenya	47
Figure 14:	Map of Lion Range in Kenya	50
Figure 15:	Map of Hyena Range in Kenya	53
Figure 16:	Map of Known Cheetah Populations	56
Figure 17:	Map of Wild Dog Range in Kenya	59
Figure 18:	Important Bird Area sites in Kenya (Source: IBA 2019)	62
Figure 19:	A Map Showing the Distribution of Birds' Sightings in Kenya Using Citizen Science (Source: Kenya Bird Map, July 2021)	63
Figure 20:	A Map Showing the Aerial Survey Area During the Tsavo Ecosystem Census, June 2021	65
Figure 21:	LSMM Ecosystem Aerial Census Blocks, June 2021	68
Figure 22:	Map Showing the location of Maasai Mara ecosystem	70
Figure 23:	A Map of the Amboseli-Magadi Ecosystem	72
Figure 24:	Athi-Kapiti Ecosystem Covered During May, 2021 Census	74
Figure 25:	Map Showing the Lamu-Lower Garissa Ecosystem	76
Figure 26:	Map of the Study Area Showing the Protected Areas and Conservancies	78
Figure 27:	Location of the Study Area and Isiolo Census Blocks	80
Figure 28:	A Map of LNNP used During the June 2021 Wildlife Census	82
Figure 29:	Map Showing study area of Nairobi National Park	84
Figure 30:	Map Showing the Location of Oldonyo Sabuk National Park, Machakos County, Kenya	86
Figure 31:	Location of Mwea National Reserve and Basic Infrastructure within the Reserve	88
Figure 32:	Map of Ruma National Park Showing Flight Path	90
Figure 33:	Study Area Shimba Hills National Reserve	92
Figure 34:	Map of the Forest Patches along the Lower Tana River Extending from Nkanjoja to Onkolde Indicating the Location of TRNPR and Including the Un-censused	94
Figure 35:	A map of Mandera County Showing the Constituencies	96
Figure 36:	A Map of Garissa County Showing the Constituencies	98
Figure 37:	A Map of Wajir County Showing the Location of the Constituencies	100
Figure 38:	Map Showing the Location of Turkana	102
Figure 39:	Map Showing the Location of Marine	104
Figure 40:	Map of the Study Area Showing the Extent, County Boundaries, and the KWS Conservations Regions.	107

ABBREVIATIONS

AED	African Elephant Database
AME	Ambosemi-Magadi Ecosystem
ANP	Ambosemi National Park
ASALS	Arid and Semi Arid Lands
CBD	Convention of Biological Diversity
CCA	Coast Conservation Area
CITES	Convention on International Trade of Endangered Species of Wild Fauna and Flora
CMS	Convention on the Conservation of Migratory Species of Wild Animals
DRSRS	Department of Resource Censuss and Remote Sensing
ECA	Eastern Conservation Area
GEC	Great Elephant Census
GoK	Government of Kenya
HCP	Hirola Conservation Programme
HEC	Human Elephant Conflict
IUCN	International Union for Conservation of Nature
KFS	Kenya Forest Service
KWCA	Kenya Wildlife Conservancies Association
KWS	Kenya Wildlife Service
LNNP	Lake Nakuru National Park
LSMM	Laikipia-Samburu-Marsabit-Meru
MCA	Mountain Conservation Area
MIKE	Monitoring the Illegal Killing of Elephants
MNP	Meru National Park
MT&W	Ministry of Tourism and Wildlife
NCA	Northern Conservation Area
NGO	Non-Governmental Organization
NMK	National Museums of Kenya
NNP	Nairobi National Park
NRT	Nothern Rangeland Trust
PA	Protected Area
SCA	Southern Conservation Area
SHIFOGA	Shimba Hills Forest Guides Association
SHNR	Shimba Hills National Reserve
TCA	Tsavo Conservation Area
TENP	Tsavo East National Park
TWNP	Tsavo West National Park
WCA	Western Conservation Area
WCMA	The Wildlife Conservation and Management Act, 2013
WCS	Wildlife Conservation Sociey
WRTI	Wildlife Research & Training Institute

EXECUTIVE SUMMARY

Kenya has a rich wildlife resource which is the main tourism product, a key economic pillar of vision 2030. However, the country is not receiving maximum benefits from the wildlife resource due to inadequate knowledge on the status and distribution of the country's wildlife populations as censuses are ecosystem and species based with no comprehensive nationally coordinated wildlife census undertaken.

The Wildlife Conservation and Management Act (WCMA) - 2013 requires the Cabinet Secretary to submit and report biannually to the National Assembly a National Wildlife Conservation Status Report as well as provide the status of wildlife monitoring every five years.

The Ministry of Tourism and Wildlife (MoT&W), the Kenya Wildlife Service (KWS) and the Wildlife Research and Training Institute (WRTI) undertook the national wildlife census for a period of 3 months (April-July 2021). The specific objective of the national census was to establish a National baseline of wildlife species. The census targeted terrestrial and aquatic ecosystems. The census enumerated land and water mammals, key birds and endangered primates. This was achieved through total aerial counts, sample aerial counts, ground counts, individual identification and indirect methods such as dung counts and use of camera traps and desktop review of previous census. The census involved the use of aircrafts (fixed-winged and helicopters), boats, vehicles, dictaphones, cameras, camera traps and used Geographic Information Systems (GIS) to record and analyse observations.

The census was officially launched by the Cabinet Secretary, Ministry of Tourism and Wildlife Hon. Najib Balala on the 6th May 2021 at Shimba Hills National Reserve. A total of 100 personnel were involved in various census sites and were drawn from Government Agencies at National and County levels, conservation partners, private and community conservancies, local communities among other stakeholders.

The census covered over 343,380km² (total aerial = 186,296km² and sample aerial = 157,084km²). This represents 59% of Kenya's land mass. A total of 1158.23 hours was flown (940.80 hours and 217.43 hours for total and sample aerial censuses respectively).

Over 30 species of mammals, birds and marine species were counted in various ecosystems during the census. Some of the iconic species counted were elephants (n=36,280), black rhino (n=897), white rhino (n=842), northern rhino (n=2), lions (n=2,589), hyenas (n=5,189), Cheetahs (n=1,160), wild dogs (n=865), buffalo (n=41,659), Maasai giraffe (n=13,530), reticulated giraffe (n=19,725), Nubian's giraffe (n=938), common zebra (n=121,911), Grevy's zebra (n=2,649), eland (n=13,581), hartebeest (n=7,332), wildebeest (n=57,813), Grant's gazelle (n=66,709), Thomson's gazelle (n=18,277), hirola (n=497), topi (n=8,627), impala (n=25,541), oryx (n=11,765), kudu (n=2,524), gerenuk (n=13,400), waterbuck (n=1,071), warthog (n=8,013), hippo (n=1,788), Maasai Ostrich (n=2,354), Somali Ostrich (n=6,483), roan antelope (n=15), sable antelope (n=51), Mountain bongo (n=150), Sitatunga (n=473), Tana river Mangabey monkey (n=1,650) and Tana river red colobus monkey (n=1,219), among other terrestrial species.

Key marine species counted were, sea turtle nests (n=340), dugong (n=2), whale shark (n=9), Humpback whales (n=63), Indo-Pacific bottlenose dolphin (n=354), Blue whale (n=2), Scalloped Hammerhead shark (n=222), Silky Shark (n=495), Giant Guitarfish (n=89), Bowmouth guitarfish (23), Mobula rays (n=11), Oceanic whitetip shark (n=358), Long comb sawfish, narrow snout sawfish or green sawfish (n=60), Great white shark (n=29) and Tiger shark (17) among other marine species.

Among the water fowl birds counted were: the lesser flamingoes (n=97,8005 individuals), greater flamingoes (n=748 individuals), great white pelicans (n=523), pink-backed pelicans (n=360), great comorants (n=4,292 individuals), long-tailed comorants (n=733 individuals), African darters (n=639), African fish eater (n=57 individuals), yellow-billed stork (n=720 individuals), Egyptian geese (n=963) and red-knobbed coot (n=692 individuals).

The census recorded an increase on some of the charismatic species such as: elephants, rhinos, lions, giraffe, Grevy's zebra, hirola among others and established baseline number for other species. However, there was relatively lower records of the plain game species.

A number of anthropogenic activities such as livestock incursions, logging, charcoal burning, settlements and fires were observed in conservation areas.

The report identifies some of the challenges such as habitat loss, land use and land tenure systems change, exponential human population growth, and accompanying demand for land for settlement and infrastructure as the reasons for low densities of plain game.

This report gives an overview of the national wildlife census results. More detailed technical reports for each ecosystem can be availed on request from Wildlife Research and Training Institute (WRTI).

Results from this national census has provided important baseline data to inform policy to sustainably conserve and manage Kenya's wildlife resources for current generations and for posterity and propose the following recommendations:

1. The National Wildlife Census to be conducted after every three years in line with wildlife ecological cycles. In this regard, a budgetary allocation by the National Treasury during the budget cycle every three years is critical.
2. An annual budget is required to support annual and active monitoring of rare and endemic species such as roan antelope, sable antelope, mountain bongo, rhino, Grevy's zebra, hirola and sitatunga
3. An immediate budget support is required to establish status of species currently threatened by illegal trade and bush meat such as pangolins, dik diks and gazelles and those species that were not covered during this census such as leopard, small carnivores and non-human primates
4. There is need for review of legislation to recognize community conservancies as protected areas as they constitute important wildlife range
5. To improve the ecological integrity of dormant parks and reserves, there is need to enhance public-private partnerships
6. A budgetary support is required to establish a data portal and access mechanism at the Wildlife Research and Training Institute to inform wildlife management as part of its mandate
7. For the endangered, rare and endemic species such the black rhino, mountain bongo, roan antelope, sable antelope among others, there is need for budgetary allocation to support development and implementation of their recovery plans.
8. Deliberate efforts to secure existing rhino population and additional space for rhinos due to the huge investment costs, security and management requirements
9. Invest in new innovations, modern census equipment and software to improve efficiency in undertaking wildlife censuses.
10. The County Governments to incorporate the census findings in the development of the County Integrated Development Plans (CIDP) and spatial plans.
11. The WRTI to pilot models of wildlife utilization programme as part of management of common species to enhance benefits to land owners who live with wildlife.
12. Fast track the full operationalisation of the WRTI which is the corporate body mandated to undertake and coordinate wildlife research.
13. Undertake a national classification of species with low populations in consultation with IUCN.

Keywords: *Wildlife Census, Aerial Total Count, Aerial Sample Count, Ground Count, Protected Areas, Marine Ecosystem, Terrestrial Ecosystems, Community Conservancies*



Message From The President

Kenya's heritage of natural beauty and scenic splendour is made infinitely richer because of our diverse wildlife. Across the globe, our Nation is renowned for our rich bio-diversity that spans air, water, and land; natural wealth that makes us a conservation and tourism destination second to none.

As the world grapples with both climate change and human activities that threaten botanical and zoological life as well as their habitats, Kenya is leading the way by implementing bold and decisive actions to conserve and promote our ecological wealth.

It is against that backdrop that the inaugural nationwide wildlife census has been conducted. This national endeavour is a historic opportunity to develop an annotated inventory of our wildlife resources, in recognition of their place as strategic national assets. The report arising from this noble national effort provides the required information to guide future conservation and management of our wildlife resources, in a manner that minimizes human-wildlife conflict and also promotes sustainable development.

As we continue to uphold our sacred duty of protecting our wildlife for current and future generations, I call on all players within the Government as well as in our private sector to make this report a central cog of all plans and actions within wildlife conservation and management.

Uhuru Kenyatta, CGH
President of the Republic of Kenya and
Commander-In-Chief of the Defence Forces



Foreword

The National Wildlife Census 2021 is a proud undertaking by the Kenya Government. It is indeed a historical achievement of documenting the first comprehensive report on the status of our wildlife as one data layer. This milestone is also a noteworthy reflection of the potential of collaboration and testimony to the capacity of our local institutions. This inaugural undertaking was coordinated by the Kenya Wildlife Service and the newly established Wildlife Research and Training Institute, who will be the repository for the National wildlife conservation status data.

The Ministry is required to report biannually to the National Assembly through the National Wildlife Conservation Status Report on the status of efforts to develop and implement recovery plans for all nationally listed species and on the status of all species for which such plans have been developed.

In the past, Kenya in collaboration with partners has undertaken site or species specific wildlife censuses. Markedly, this is the first one-off national wildlife census that has been carried out aimed at establishing the most comprehensive baseline on wildlife status and distribution in the country.

Obtaining this level of information of a vital natural resource allows for better policy, planning and assessment of areas that require focus in our interventions to maintain or improve our National conservation efforts. This also guides the strategic deployment of vital resources aimed at increasing numbers where declines are evident as well as drawing attention to areas that require mitigation measures to avert potential or real human-wildlife conflicts and eradicating the threats posed by poaching and illegal bush meat trade.

The findings of this report will also provide a scientific basis for programmes dedicated to saving our endangered species and their habitats for future generations.

Evidence from the census illustrates the social-economic impacts of activities such as agriculture, human settlements and infrastructure development on wildlife movements and loss of space for wildlife. This will require more attention if we are to avert the danger of secluding wildlife in pockets of Protected Areas. The world-renowned success of the Kenya's model of free-ranging wildlife is based on our ability to allow as much unhindered movement and distribution of wildlife. It is imperative that we continue to strive to maintain this model while taking into account the demands of a growing human population and the need for clear benefits.

It is important to recognize and appreciate that maintaining this successful model has a huge cost in terms of both finances required and impact on humans. It is therefore vital that the global community takes this cost as a shared responsibility. This calls for increased investment by development partners and also by the private sector in taking up a bigger role in the form of Public-Private Partnerships in biodiversity conservation and protected areas.

We conserve this magnificent heritage for the posterity of Kenyans and the entire World.

Hon. Najib Balala, EGH

Cabinet Secretary, Ministry of Tourism and Wildlife



Preface

Kenya is developing at a fast rate with new and major infrastructural projects (road networks, railway lines and urban centres) being completed or in the process of being developed. Also, emerging incompatible changes in land use and land tenure systems, especially in rural areas that are important wildlife ranges, have serious implications for wildlife numbers and distribution.

These are exacerbated by the ever increasing human population that increases demand for shared and limited resources. The areas have also been experiencing erratic weather conditions resulting in droughts as global climate change impacts are felt in Kenya. This is likely to impact negatively on wildlife population for certain species. Understanding wildlife population status, trends and distribution in relation to developments is key to initiating species specific and habitat conservation actions. Therefore, this national wildlife census has provided information on the wildlife numbers, their trends, distribution and threats to guide development of such conservation actions. In addition, the information will be used to guide policy and decision making processes as well as advice on the appropriate management actions required to ensure stable and increasing wildlife populations including expansion of wildlife range.

The national census results will also enhance understanding of the connectedness of the key landscapes. For example, past monitoring has shown extensive cross landscape wildlife movements between the Amboseli and Tsavo Ecosystem, Meru and Laikipia ecosystem as well as Laikipia and Marsabit ecosystem. Isolated populations reduce genetic diversity and health of ecosystems and therefore connectivity mitigate problems associated with fragmentation by joining isolated wildlife populations. It is expected that by having a one-off census, clear understanding of national and ecosystems wildlife population status and distribution will be understood. The data will form the basis for future wildlife population monitoring as well as establishing real economic value of wildlife capital for reflections in the national budgeting process.

Peer reviewed scientific methods for wildlife censuses for different target wildlife species were used to gather and provide data used in this report. The report provides population status, trends and distribution of different wildlife species, which include elephants, rhino, sable antelope, roan antelope, buffalo, giraffe, Grevy's zebra, common zebra, giraffe, eland, topi, kudu, impala, hirola, gerenuk, warthog, hippo, ostrich, crested mangabey, red colobus monkey, sitatunga, oryx Grant's gazelle, Thomson's gazelle.

This report will guide efforts toward reversing the declining wildlife population as well as develop strategies to grow stagnant populations. The implementation of outcomes from this report and future one-off censuses will require financial and human resources support. Therefore, I call upon our development partners, conservation NGOs and other stakeholders to support the WRTI and the KWS to ensure the implementation of the outcomes and future censuses are realized.

PROF. FRED H. K. SEGOR, CBS

Principal Secretary, State Department for Wildlife

Acknowledgement



Brig. (Rtd) John M. Waweru, EBS
Director General
Kenya Wildlife Service



Dr. Patrick Omondi, OGW
Ag. Director/CEO
Wildlife Research and Training
Institute

The Country undertook for the first time a one-off national wildlife census to establish the status of her wildlife resources to facilitate planning for the resources. Successful implementation of the National Wildlife Census with the slogan #CountToConserve# was undertaken with budgetary allocation from the Government through the National Treasury. We thank HIS EXCELLENCY HON. UHURU KENYATTA, C.G.H., PRESIDENT OF THE REPUBLIC OF KENYA AND COMMANDER IN CHIEF OF THE DEFENCE FORCES for prioritizing the wildlife census and allocating financial resources to undertake the exercise despite the tight budgetary constraints to various sectors amidst impacts of the COVID-19 pandemic.

We thank The Cabinet Secretary Hon. Ambassador Ukur Yatani, EGH Cabinet Secretary, National Treasury & the Ministry of Planning for the timely dispatch of funds to facilitate the census.

We also thank the Cabinet Secretary, Ministry of Tourism and Wildlife Hon. Najib Balala, CBS for policy guidance on the entire process and officially launching the National Wildlife Census in Shimba Hills National Reserve on 6th May 2021, The Chief Administrative Secretary, Ministry of Tourism and Wildlife Mr. Joseph Boinet, the Principal Secretary, State Department of Wildlife, Ministry of Tourism and Wildlife (Prof. Fred Segor) and staff of Ministry of Tourism and Wildlife for their technical and logistical support. The Ministry of Interior and Coordination of Government is highly appreciated as the County Commissioners ensured smooth operation of the teams by ensuring the teams' security and informing communities about the census through the elaborate networks of administrators across the Country up to the village level. All other Government Ministries who provided support to facilitate implementation of the census are also appreciated in particular the Kenya Defense Forces, National Air Support Department (NASD) for aviation support and logistics, Department of Resource Census and Remote Sensing, Kenya Forest Service, Kenya Maritime Authority, State Department for Fisheries, Kenya Marine Fisheries Research Institute, Kenya Coast Guards, National Museums of Kenya, Institute of Primate Research, Lake Victoria Basin Authority and Community Scouts under the Government economic stimulus package.

The County Governments are thanked for facilitating the census in their respective counties whose outcome will be key in informing County spatial planning process and inform areas of collaboration.

We thank the Conservation Alliance of Kenya, particularly the following Non-Governmental Organization (NGOs), Conservancies and Individuals: World Wildlife Fund(WWF) for Nature- Kenya, Africa Wildlife Foundation(AWF), Save The


Elephants(STE), Space for Giants(SFG), Mara Elephant Project(MEP), Amboseli Elephant Trust (AEP), Tsavo Trust, Sheldrick Wildlife Trust, Big Life Foundation, Amboseli Ecosystem Trust, Wildlife Conservation Society, Coastal Oceans Research and Development in the Indian Ocean, Arocha Kenya, Nature Kenya, Tropical Sea Life, Community Based Environmental Conservation, Olive Ridley, Maasai Wilderness Trust, Stand Up Shout Out, Lewa Conservancy, Loisaba Conservancy, Olpajeta Conservancy and Northern Rangeland Trust and Maasai Mara Wildlife Conservancies Association Captain Wangai Kimeria, Captain Justin Mathews and Captain Mark Ross for supporting the census with aircrafts and personnel. Kenya Wildlife Conservancies Association and its membership are also appreciated. We also thank Community Based Organizations among them Watamu Turtle Watch, Diani Turtle Watch, TIWI Turtle Watch Police, Beach management units, Friend of Arabuko Soke Forest and Shimba Hills Forest Guides Association.

Institutions of higher learning, research institutions and projects are also appreciated in facilitating and participating in the census in particular Technical University of Mombasa, Kenyatta University (Mombasa Campus) and Mpala Research Centre.

The media fraternity and the census digital and documentary team is acknowledged for documenting and disseminating information to both the wider public and international community and for putting together digital records for future reference. Special thanks go to Digimatt Solutions and Dharkemmy Corporate Communications Limited (DCCL) for their cooperation in this regard.

Finally, special thanks go to the National Wildlife Census Planning Committee, Kenya Wildlife Service and Wildlife Research and Training Institute staff for ensuring articulate planning, organization and implementation of the census. The team that prepared the report is also acknowledged for working round the clock to ensure the national census report was finalized and submitted within the required timelines.





CHAPTER

1

General Introduction



1.1 | BACKGROUND

Kenya's rich wildlife resource is one of the key economic pillars of the country. Wildlife-based tourism is a cultural service that is an important source of revenue accounting for about 10% of the Gross Domestic Product (GDP) of Kenya in 2010 (Kibara et al., 2012; GoK, 2010). However, the country has not been receiving optimized wildlife-based benefits as anticipated due to inadequate knowledge on the status of the country's wildlife resources as well as factors influencing their abundance and distribution. This is because there has been no one-off and comprehensive scientific data on wildlife population to inform sound policy direction. To achieve this, the National Government provided funding to facilitate a coordinated national wildlife census in all regions that harbour wildlife including protected areas, conservancies, private and community areas. The information generated will support development and implementation of sound wildlife conservation policies. In addition, the information will be used to guide evidence-based decision support tools for adaptive wildlife management, integrated cross sectoral and wildlife conservation, multi-scale planning and sustainable development to optimize wildlife benefits to the country. This will be more so to the wildlife species listed in schedule six (6) of the Wildlife Conservation and Management Act (WCMA) - 2013 hereafter referred to as WCMA-2013.

The number and distribution of species listed in schedule six (6) of the WCMA - 2013, (i.e., rare, endangered and threatened species) whose population status and distribution has not been evaluated recently require urgent update. Such updates on these species population status and distribution can be achieved using different methods. These include; aerial total census, aerial sample census, ground census using vehicles and indirect methods like the dung count and use of camera traps (Norton-Griffiths, 1978; Douglas-Hamilton, 1996). Aerial censuses are commonly undertaken in savannahs, arid and semi-arid ecosystems and requires use of fixed wing light aircrafts and helicopters. Within the forested ecosystems, the use of indirect dung count methods and camera traps targeting elusive forest species are commonly adopted.

After the enactment of the WCMA-2013, wildlife censuses become a mandatory legal requirement. Section 64(3) of the WCMA 2013 stipulates that the Cabinet Secretary (CS) responsible for wildlife matters is required at least once every five years, submit to the National Assembly (NA) a wildlife resources monitoring report showing the

achievements made in the implementation of the past or subsisting national wildlife conservation and management strategy and avail the said wildlife resources monitoring report to the public. Therefore, this Government initiative to provide financial support to undertake the National Wildlife Census aims to fulfil the legal requirement as stipulated by the WCMA - 2013. Previous censuses were undertaken using both Government funds as well as from conservation partners and Non-Governmental Organizations (NGO).

Aerial total censuses have been carried out over the years in the larger ecosystem including the Tsavo, Amboseli, Maasai Mara and Laikipia-Samburu-Marsabit extending to the Meru Conservation Area (MCA) in 2002, 2005, 2006, 2007, 2011, 2014, 2015, 2017 and 2018 (Omondi et al., 2002; Chase et al., 2015; Thouless et al., 2016; Ngene et al., 2017; Kyale et al., 2018; Ngene et al., 2018). These regular censuses have become a manager's tool to determine wildlife trends, establish national wildlife databases, manipulate the range, condition, productivity, and the species numbers, including distribution and provision of space and grazing. Furthermore, security and tourism are also influenced by the numbers within and outside protected areas. Besides, human-wildlife conflict issues are also adequately addressed as well as resource allocation towards conservation and monitoring of the same.

Kenya has never undertaken a one-off national census to establish a baseline of the wildlife population status and distribution across country. It was therefore important to undertake this national census to gather the baseline data on the current wildlife population status and distribution as well as future use to understand wildlife population trends and shifts in their distribution. During the census, terrestrial, fresh water, marine mammals, key birds (ostrich and kori bastards), endangered primates (Tana Mangabey and Tana red colobus) and reptiles (crocodiles) were counted. Different scientific methods were applied depending on the species and the habitat types following standard methods. Aerial census techniques were used to count large mammals in savannah, arid and semi-arid regions while dung count methods were used in forested ecosystems. For the wetland ecosystems, both aerial and ground count methods were applied. Existing data of other species of wildlife was collated from different conservation institutions and synthesized to produce this census report. This report presents the finding of the first ever National Wildlife Census (NWS) in Kenya.

1.2 | GOAL AND OBJECTIVES

The goal of this aerial total and sample censuses was to establish a national baseline of wet season wildlife population status and distribution in Kenya. The specific objectives of the National Wildlife Census were to:

- a. Determine Kenya's wildlife population abundance and distribution
- b. Identify threats to wildlife conservation and management in Kenya's landscapes
- c. Establish the suitability of use of drones in aerial censuses in open savanna ecosystems
- d. Establish a national wildlife database and data portal
- e. Establish resource capacity for undertaking wildlife censuses
- f. Suggest strategies for effective wildlife conservation and management across Kenya's landscapes.
- g. Determine Kenya's wildlife population trends over time
- h. Provide data and information for use to prepare a national wildlife status report

1.3 | PROJECT JUSTIFICATION

Kenya's human population has grown 5.95-fold from 8 million people in 1963 to 47.6 million in 2019, however the available surface land remains the same. Increased human population has led to increase in demand for land for human settlements, infrastructure, and agriculture. This has resulted in land use changes and tenureship which have important implications for sustainable wildlife conservation in this country. Competing and non-compatible land uses threaten biodiversity conservation in Kenya, and therefore understanding wildlife abundance and distribution in relation to these developments is key to initiating habitat conservation actions.

Key wildlife landscapes in Kenya have been experiencing challenges in terms of land tenure, land use system, increased livestock and poor husbandry practices, reduced rainfalls and increased droughts occasioned by global climate change amongst others. This is likely to

impact negatively on certain wildlife species populations. This national census is expected to provide information on the wildlife numbers and trends in all the ecosystems, and therefore propose appropriate management actions.

The census enhanced understanding of the connectivity of the key landscapes. For example, past monitoring has shown extensive cross landscape wildlife movements between the Amboseli Ecosystem and Tsavo Ecosystem, Meru Ecosystem and Laikipia Ecosystem as well as Laikipia Ecosystem and Marsabit Ecosystem. It is expected that by having a one-off census, clear understanding of national wildlife population status and distribution will be understood. The data forms the basis for future wildlife population monitoring. In addition, the data forms a basis for establishing real economic value of wildlife capital for reflections in the national budgeting process.





CHAPTER

2

Materials and Methods



2.1 | STUDY AREAS

The census was carried out in the following ecosystems, protected areas, wetland areas and counties: Tsavo ecosystem; Amboseli-Magadi ecosystem; Athi-Kapiti ecosystem; Naivasha-Nakuru ranches ecosystem; Laikipia-Samburu-Marsabit-Meru ecosystem; Nasalot-South Turkana-Rimoi ecosystem; Shimba Hills-Mwaluganje ecosystem; Lamu - Lower Garissa and Tana River ecosystem, Mwea National Reserve; Ruma National Park; Lake Nakuru National Park; Nairobi National Park; Rift

Valley Lakes; Sitatunga range in wetlands within Western and Central Rift Conservation Areas; Garissa County; Wajir County; Mandera County; and, Turkana County. Together, these areas constitute some 343,380 km². These sites correspond with the Kenya Wildlife Service (KWS) Conservation Area Network. Figure 1 below shows the census areas in Kenya. Detailed description of each census region is provided in the separate reports of each census site.




Figure 1: A map showing the Kenya Wildlife Service Conservation Area where the census took place between April and July 2021

2.2 | METHODS

The following standard methods were used during the National Wildlife Census to count different species of wildlife in different conservation areas:

1. Aerial total census method was used in Tsavo, Laikipia-Samburu-Marsabit-Meru-, Isiolo, Amboseli-Magadi, Athi-Kapiti, Maasai Mara, Lamu - lower Garissa and Tana River, Kerio Valley and Nasolot ecosystems; Rift Valley lakes; Ruma, Sibilo, and Lake Nakuru National Parks; Mwea National Reserves, and Shimba Hills National Reserve and Mwalungaje Sanctuary; and Solio Ranch following procedures described in details by Norton-Griffiths (1978) and Douglas-Hamilton et al. (1996).
2. Elephants in forested ecosystems of Aberdare National Park, Kipipiri Forest Reserve, Mount Kenya National Park & Reserve, Mount Elgon National Park, Chepkitala Forest Reserve, Shimba Hills National Reserve, Kiriya Forest, and Loita Forest were counted using the indirect dung count method as described by Barnes et al. (1997) and modified by Vanleueew (2008).
3. Censuses for rhino meta-populations hosted in 19 smaller populations in public, private, county and community areas were undertaken through ground, aerial, water hole census and camera traps. Detailed methods are describe in the separate reports on population status and distribution of rhinos in Kenya.
4. Ground census was carried out to establish the population of roan antelope in Ruma National Park, sable antelope in Shimba Hills National Reserve, endangered primates within Tana River forest complex ecosystem. Walking transects were used to count the roan antelope, sable antelope, red colobus and the crested mangabey. In small protected areas like Nairobi and Oldonyo Sabuk National Parks, transect counts using vehicles and on foot respectively were undertaken to establish the population status and distribution in the two protected areas. Wetland birds were also counted using standard ground census methods. All the ground census methods followed standard procedure as outlined in details by Norton-Griffiths (1978). Detailed methods are described in respective chapters outlined in this report.
5. Censuses for wildlife species within wetlands and lakes (e.g., crocodiles, hippos and waterfowl birds) was carried out along transects using aircrafts and boats following standard methods outlined in details by Kanga et al.(2019).
6. The exercise also involved digitization of all collected scientific data, preparation and compilation of publications and databases.
7. The data was summarized using Argis 10.0 as described by ESRI (2010) and analyzed following procedures described by Zar (1996).
8. Drone technology was piloted in 5 sites of Tsavo, Mwea, Nairobi, Nakuru and Ruma, the results of the pilot to inform future applicability.

hughran.







CHAPTER






3






2021 Wildlife Census Results Summary
























WILDLIFE NUMBERS AT A GLANCE












3.2 | KEY WILDLIFE SPECIES





AFRICAN ELEPHANT

Loxodonta africana



DID YOU **KNOW?**



Elephants have around **150,000** muscle units in their trunk.

Their trunks are perhaps the **most sensitive organ** found in any mammal.



UPTO **3.2M**



UPTO **6000KG**



EN

3.1.1 Elephant

Due to the charismatic nature of the African elephant, and its ability to modify its natural habitats, hence affecting the occurrence of other species where they range, monitoring elephant for provision of accurate and timely information on their population status is paramount. The Government of Kenya (GoK) through its arm Kenya Wildlife Service (KWS) is obligated to carry this function, in collaboration with relevant conservation partners.

Dispersal is a critical component of animal ecology that is poorly understood for most species. In particular, savanna elephants (*Loxodonta africana*) have been studied for decades in national parks across Africa, but little is known about their dispersal into new or unused habitats or their population dynamics in human dominated landscapes. This is exacerbated by the immiserating community livelihoods around elephant range areas.

Continued monitoring has led to the success in the identification and response to poaching and Human-Elephant Conflict (HEC). The monitoring has been facilitated by improvement on monitoring technologies and methodologies. Detailed genetic analysis of the elephant population has resulted to the recent confirmation of two distinct species of elephants, that is the forest elephant (*Loxodonta cyclotis*) and the savanna elephant (*Loxodonta africana*) (Hart et al. 2021). As aforementioned, the Kenya elephant population is therefore classified as

The Tsavo ecosystem accounts for more than 37% of the national elephant range, which is estimated at 49,000 km², followed by the Samburu-Laikipia ecosystem with 37,937 km². The Mara and Amboseli West Kilimanjaro ecosystems accounting for 11,681 km² and 37,937 km² respectively (Figure 2).






Figure 2: The Kenya National elephant range as at July 2021



RHINOCEROS

Diceros bicornis michaeli (BR)

Ceratotherium simum (SWR)

Ceratotherium cotoni (NWR)

2

CR

Northern White Rhino(NWR)



897

CR

Black Rhino(BR)

840

EN

Southern White Rhino(SWR)



DID YOU KNOW?



Black rhino has more or less the same coloration as white rhinos. Instead of the color, white rhinos are named after the Afrikaan word **wyd**, which describes the animal's **square upper lip**. Black rhinos have a **pointed upper lip**, named after the opposite color to easily differentiate them from white rhinos.

(Source: facts.net)



UPTO 1.3M



UPTO 1000KG



EN

CR

3.1.2 Rhino

The future of the black rhino is of critical importance to the Government of Kenya being a species of conservation concern following a dramatic reduction in their population in the 1970s and early 1980s from approximately 20,000 as a result of the illegal trade in their horn to fewer than 400 animals in 1987.

Kenya has the 3rd largest population of rhinos in Africa after South Africa and Namibia having a total rhino population of 1,605 (853 black rhinos, 750 southern white rhinos and 2 northern white rhinos) as at end of 2020. Kenya is home to the world's third largest black rhino population after South Africa and Namibia.

Kenya is implementing the 6th edition of the Black Rhino Action Plan (2017-2021) whose long term vision is: 'To have a meta-population of at least 2,000 black rhinos of the eastern African subspecies (*Diceros bicornis michaeli*) in Kenya, and in suitable habitats as a global heritage'. The overall goal is: 'To achieve a meta-population of 830 black rhinos by the end of 2021.

Kenya has already surpassed its goal in the Action Plan having achieved a population of 853 black rhinos as at 31st December 2020. The Black Rhino Action Plan which expires at end of 2021 is being reviewed for the period 2022-2026 with revised goals and activities.

Kenya remains the stronghold of the eastern black rhino subspecies conserving just over three quarters

(approximately 80%) of the wild population of the sub species. Kenya's meta-population has fifteen black rhino populations out of which 12 are rated as follows; four IUCN SSC African Rhino Specialist Group (AfRSG) rated Key 1 (>100 black rhinos), four Key 2 and four Important 1 populations that are of continental significance. Kenya's black rhinos are conserved in nine state, four private, one county and one community lands across the country

Kenya is for the first time developing the first White Rhino (*Ceratotherium simum*) Conservation and Management Action Plan (2021-2025) with the population of the southern white rhino having undergone rapid growth since introduction with current population estimated at 750 individuals from the 51 individuals introduced from Southern Africa in the 1960s, 1970s and 1990s thus the need to develop a separate strategy for the species to guide on best practice for conservation and management of the species.

Kenya rhino population is managed as a meta-population (A number of sub-populations of a species or sub-species managed collectively as one single population with occasional movement of animals from one sub-population to another) hosted in 17 rhino conservation sites spread across the country. Out of the 17 sites, 9 are in parks, 4 on private land, 3 on community land and





Figure 3: Map showing rhino conservation areas

AFRICAN BUFFALO

Syncerus caffer



41,659

GESTATION PERIOD
11.5 MONTHS



DID YOU **KNOW?**



African Buffalo will often kneel down and rub their necks and heads on the ground in a display of aggression to establish dominance.



UPTO 1.6M



UPTO 800KG



LC

(Source: <https://drewsproule.com/>)

3.1.3 Buffalo


Introduction

Buffalo is one of the most abundant and widely distributed species in Kenya and among the big five iconic species. According to IUCN red list, the species is considered as of no concern. However in Kenya it's a species of interest because of conflicts with human through crop raiding, human attacks and disease transmission to livestock.

It occurs in both forested and open savannah ecosystems. During this census, only those in open Savannah ecosystem were counted using aerial total count method in 14 sites. Those in forested ecosystem were not counted as a different technique is applied to assess their population which requires more time. For example the use of camera traps.

Population status of buffalo

The population of buffalo counted in the open savanna ecosystems during the current national census were 41,659 animals. They were recorded in 14 sites (Table 9). The Maasai Mara ecosystem recorded the highest population of buffalo in the country (n=11, 604 buffaloes), which represents about 28% of the total buffalo population in Kenya. This is followed by Tsavo ecosystem (19%), Lake Nakuru National Park (15%) and then Laikipia-Samburu-Marsabit ecosystem (15%). These four ecosystems account for about 78% of the total Kenya's buffalo population. Small isolated populations of buffalo were observed in Ol Donyo Sabuk NP (23), Shimba Hills NR (23), Athi-Kapiti ecosystem (30) and Mwea National Reserve (69).






Figure 4: The distribution of buffalo in Kenya (2021)

GIRAFFE

G. tippeiskirchi (Masai Giraffe)

G. reticulate (Reticulated Giraffe)

G.c. camelopardalis (Rothschild's Giraffe)

768 EN

Nubian Giraffe

13,732 LC

Maasai Giraffe

19740 LC

Reticulated Giraffe



DID YOU KNOW?



A giraffe's spots are like human fingerprints.

Aside from the fact that giraffes have seven vertebrae in the neck like humans, its spots in the body are much like human fingerprints.



Masai G.

Reticulated G

Rothschild's G.



UPTO 5.7M



KG

UPTO 1200KG



LC

EN

3.1.4 Giraffe

Kenya is home to three species of giraffe, which include the Maasai giraffe, Reticulated giraffe and the Nubian giraffe. Their total population estimate is 34,240 animals (Maasai giraffe = 13,530 individuals; reticulated giraffe = 19,740 individuals; and, Nubian giraffe = 970 individuals. Table 10 below provides a summary of the population status of the three species in Kenya as at July 2021.

The three giraffes have distinct geographical location (figure 11) with Maasai giraffes occurring mainly in Kajiado, Narok, Makueni and Taita Taveta Counties. Reticulated giraffe were observed in the North-Eastern counties of Wajir, Garissa, Mandera, Turkana including Laikipia, Samburu, Lamu and Tana river while Nubian's giraffe was observed

mostly in Ruma National Park in Homa Bay county, Lake Nakuru National Park and Naivasha ranches in Nakuru county, Mwea national reserve in Embu County and Nasalot & Kerio valley in Baringo/Elgeyo Marakwet counties. Within captive facilities, a substantial population of 47 individuals of all species is conserved in different private facilities in Kenya.

The key stronghold for Maasai giraffe is Amboseli (n=6,425) and Tsavo ecosystem (n=4,314) while Reticulated giraffe is mostly found in Wajir, Garissa and Lamu Counties. Over 56.7% (n=550) of the Nubian giraffes in Kenya were observed in Ruma National Park.







Figure 5: Distribution of Giraffes in Kenya

GREVY'S ZEBRA

Equus grevyi



2,649

GESTATION PERIOD



12 MONTHS

DID YOU **KNOW?**



Unlike the other zebras, Grevy's Zebra, also known as Imperial Zebra it is tall, its stripes are narrower, and has big ears. It can live up to five days without water.



UPTO 1.6M



UPTO 450KG



EN

(Source: facts.net)

3.1.5 Grevy's Zebra

The national grevy population estimates are displayed in Table 11 below. A total of 2,649 grevy zebras were counted with the majority (38 %) found in Laikipia County followed by Isiolo County with 703 individuals (27%). The least number were directly sighted in Garissa County. However,

it should be noted that the methodology used to count the grevy zebra in Wajir and Garissa counties was not the most ideal as the transects were spaced 5 Km apart. The Grevy's zebra population in Naivasha County are found in Oserian ranch.






Figure 6: Grevy's zebra distribution map in Kenya

SABLE ANTELOPE

Hippotragus niger



DID YOU KNOW?

Sable antelopes congregate in herds of 15 to 20 individuals with **matriarchal social structure**.

Within the group, the more dominant female is the leader. There is only one adult male (called a bull) in each herd. The juvenile males are exiled from the herd at about 3 years old.

(Source: <https://drewsproule.com/>)



UPTO 1.9M



UPTO 235KG



CR

3.1.6 Sable Antelope



A total of 51 sable antelopes comprising 2 males, 20 females and 29 unsexed individuals were counted in the latest Sable census (Figure 16). The population of sable established in this census could be considered the lower limit estimate since it was conducted during the wet season when the species sightings in the reserve are obviously low. This could explain the very low identification of species ending up with high number of unsexed individuals as compared to the 2016 census (Table 13) which recorded a minimum of 28 males and 58 females.




Figure 16: Sable antelopes counted in the last census in Shimba Hills ecosystem




Figure 7: Historical distribution of Roosevelt's sable antelope in Kenya from 1884 to present
(Source: Modified from Butynski et al., 2015)

MOUNTAIN BONGO

Tragelaphus eurycerus isaaci




DID YOU KNOW?



In order to swiftly maneuver through the dense forest vegetation, bongos **tilt their chin up**, causing their horns to lie flat

against their back. They take this position so frequently older bongos often have bald spots on their back from the tips of their horns rubbing away the fur.

(Source: <https://drewsproule.com/>)



3.1.7 Mountain Bongo



The Kenyan mountain bongo (*Tragelaphus eurycerus isaaci*) is a critically endangered tragelaphine antelope sub-species only found in Kenya endemic to the Aberdare, Mount Kenya, Cheranganis Hills and the Mau Forests Complex, with only a few individuals left in the Eburu, Maasai Mau and South Western Mau. The species numbers and range has undergone a drastic decline in all these forests with limited information on the exact number of animals, though inferential figures stand at less than 100 individuals mainly confined to the Aberdare and Maasai Mau (East, 1999; Reillo, 2002, unpublished report BSP 2016).

In 2003, bongo repatriation from the USA was initiated to establish a sustainable, in situ managed bongo population at the Mt. Kenya Game Ranch now referred as Mount Kenya Wildlife Conservancy (MKWC) from which multiple wild-population recovery strategies could evolve. The principal objective of this project was to establish an in situ captive breeding program, in a natural setting, as the first phase of several conservation steps required to reintroduce mountain bongos to the wild. The project aimed to re-establish a viable and self-sustaining population in the bongo's native habitat.

The mountain bongo is listed as Critically Endangered by the IUCN/SSC Antelope Specialist Group (IUCN, 2003) and listed on Appendix III of the Convention on International

Trade of Endangered Species of Flora and Fauna (CITES), which allows limited trade on the species. In Kenya, bongos are accorded full protection under the Wildlife Conservation and Management Act, 2013.

Kenya has developed the 1st National Recovery and Action Plan for The Mountain Bongo (*Tragelaphus eurycerus isaaci*) in Kenya (2019-2023) VISION: We envisage viable, free-ranging and genetically representative populations of mountain bongo, thriving across intact historic mountain ecosystem ranges, cherished by the Kenyan people and the global community.Goal: To secure minimum population size for mountain bongo within their ranges in Kenya, to achieve a national population of 730 individuals over the next 50 years

Census Methods

Mountain Bongo's are shy and skittish being found in thick forests thus difficult to be easily sighted and counted. Most of the sightings are opportunistic and use of indirect methods such as dung. Camera trap images for individual ID are used to estimate the wild populations in Masai Mau, South West Mau, Eburru, Aberdare NP and Ragati in Mount Kenya NP. For the captive population in Mount Kenya Wildlife Conservancy physical sighting and individual ID is used. The current Mountain Bongo range is shown in the map in figure 20 and population estimate is shown in Table 15

Mountain Bongo




Figure 8: Map of current Mountain Bongo Range in Kenya

The mountain bongos are currently estimated at 150 individuals with 96 in the wild and 54 in captive facility in Mt. Kenya Wildlife Conservancy (MKWC) as shown in Table 15.



ROAN ANTELOPE

Hipotragus equinnus



DID YOU KNOW?



Roan antelopes have lighter underbellies, white eyebrows and cheeks and black face which is lighter in females.

The horns are ringed and can reach 1m long in males which are slightly shorter in female and hatch backwards



UPTO 2.4M



UPTO 300KG



CR

3.1.8 Roan Antelope



The WV antelope (*Hippotragus equinus*) is a rare species, restricted to savannah woodlands of Africa. It was once widely distributed through 34 eastern, central, southern and West African countries but during the last 40 years it has declined rapidly, presumably as a result of poaching and conversion of natural land into agriculture. Currently, the species possibly remains only in 30 countries (Ansell, 1971). Studies indicate that two thirds of populations throughout Africa show a decrease, while only one third is either stable or increasing, predominantly in private reserves in South Africa. The species has been extirpated from Burundi, Eritrea, Gambia and Swaziland, whereas there are an estimated 76,000 roan antelopes globally. One-third of this is concentrated in four countries: Burkina Faso; Cameroon; Zambia; and Tanzania.

In East Africa, it was confirmed to exist in small numbers in

a few protected areas, with less than 30 in Ruma National Park in Kenya, less than 30 in Pian Upe Game Reserve in Uganda and in north-western Tanzania 260 in Burigi and Biharamulo Game Reserves (Magin and Kock, 1997). The Ugandan population was thought to have gone extinct by 2003 although a few individuals have lately been sighted in Pian Upe Game reserve, while the NW Tanzania population has been much reduced from the impacts of refugee camps after the genocide in neighbouring Rwanda. There were and are reports of roan in Akagera but these have not been confirmed. Around 5,000 may still occur in southern and central Tanzania, but it is not certain that these individuals are of the same subspecies. The current situation of *H. e. langheldi* in Rwanda and Burundi is unknown, but believed to be insecure.




Figure 9: Roan distribution in Ruma N.P



Figure 10: Roan antelope sighting in Ruma NP

SITATUNGA

Tragelaphus spekei



DID YOU KNOW?



Only the males possess horns;

which are spiral in shape, have one or two twists and are 45–92 cm long



UPTO 1.25M



UPTO 119KG



EN

3.1.9 Sitatunga Antelope



The Sitatunga (*Tragelaphus spekei*) is a specialized semi-aquatic antelope adapted to living in swampy or permanently marshy wetlands in African tropics. Sitatungas occur in tall and dense vegetation of perennial and seasonal swamps, marshy clearings within forests, riverine thickets, and mangrove swamps. This medium sized antelope is highly adapted to spending the greater part of its life in papyrus swamps.

The most physical adaptation to their marshy environment is the long, splayed hooves, which enable the animal to stand and walk on mud and floating islands of vegetation without sinking. The shaggy, water-resistant coat varies in colour among populations, but is generally greyish-brown in males and rufous-brown in females and juveniles.

The species is listed in CITES Appendix III and classified

as Least Concern in the IUCN Red List (IUCN 2017). This is attributed to species poaching and loss of habitat. The species copes well with high hunting pressure in much of its range, but is most threatened by drying out of its aquatic habitat caused by changes in hydrology.

Sitatunga has specialized habitat requirements, hence their distribution range in Kenyan wetlands is patchy and discontinuous. Most of their habitats are situated outside protected areas and are facing immense anthropogenic pressure and impacting negatively on the survival of the species. The situation is compounded by limited national comprehensive studies on the species.




Figure 11: Distribution of Sitatunga's in Western conservation Area




Figure 12: distribution of Sitatungas in Central rift conservation area

HIROLA

Beatragus hunteri



DID YOU KNOW?



Both sexes have **ringed horns which hatch backwards**, in females this can reach 61 to 102

cm while in males they are 81 to 165 cm long.



UPTO 1.25M



UPTO 118KG



CR

3.1.10 Hirola Antelope

i. Introduction



The hirola's natural range – which pre-historically probably extended all the way from the Cape to the Horn of Africa (Bunderson, 1985) – has contracted over the millennia to the point where the species now occurs largely in Garissa County Kenya (Ali et al. 2017). Currently, *Beatragus* (uniquely represented by the hirola) exists only in the Ijara and Fafi sub-counties of Garissa County, between the Tana River and the Kenya–Somalia border. An area no more than 1,500 km² and another small translocated (ex situ) population established within Kenya's Tsavo East National Park in 1963 (Hofmann, 1996; Andanje & Ottichilo, 1999; East, 1999; Butynski, 1999; Andanje, 2000a, b).

The current population estimates of hirola are from the National Wildlife Census 2021 data, The Northern Rangelands Trust (NRT) Ishagbin hirola status report 2020 and Hirola Conservation Program (HCP) hirola monitoring data 2020. The Ishagbini community conservancy scouts using the Wildlife-CoMMS monitoring system collected data from The NRT. Similarly, HCP has a network of community scouts who collect hirola-monitoring data at the community areas. The National Wildlife Census 2021 employed the Total Aerial Counts methodology to collect data on the Hirola.




Figure 13: Map showing the distribution of hirola in Kenya

AFRICAN LION

Panthera leo



DID YOU **KNOW?**



That lions are **the most sociable large cat**, living in prides of between 3 and 30

individuals.



UPTO 189KG



EN

(Source: <https://drewsproule.com/>)

3.1.11 The African Lion

i. Introduction



The lion population in Africa has been on the decline in the last 100 year with the lion population in Africa declining by 96.5 percent. According to IUCN red list, in 1900, the lion population in Africa was about one million and by the 1940s, the numbers had dropped to 500,000. In 1975, the continent had 200,000 lions, but in 1990 there were just 100,000. By the year 2000, the population had declined drastically to only 35,000. Today there are about 20,000 lions with the number continuing to drop. The African lion is CITES listed and ranked as 'Vulnerable' by the IUCN Red List (Bauer, Nowell, and Packer, 2008). Population of lions

in Kenya has been on a downward trajectory with current estimate being just over 2,000 individuals.

Kenya is currently implementing the 2nd edition of the National Recovery and Action Plan for the Lion and Spotted Hyena in Kenya (2020-2030) whose vision is to sustain viable populations of lions and spotted Hyenas in healthy ecosystems as a world heritage valued by the people of Kenya while its goal is to restore and maintain viable populations of lions, spotted Hyenas and their wild prey while minimizing conflict and maximizing value to local communities.




Figure 14: Map of lion range in Kenya

SPOTTED HYENA

Hyaenidae


5,147



DID YOU **KNOW?**



The milk of a spotted hyena is **very rich**, and **cubs can last for a few days** between feeds, unlike wild dogs and lions.



UPTO 0.92M



UPTO 90KG



EN

3.1.12 Spotted Hyena

i. Introduction



Spotted Hyenas do better in many human-dominated landscapes than lions and can thrive where lion densities are lower. The IUCN redlist assessment estimates 27,000 to 47,000 spotted Hyena occur across sub-Saharan Africa. Hyenas suffer from many misconceptions, including associations with witchcraft in many parts of their range, which often make them feared or disliked by people. Largest known populations in Serengeti (Tanzania, Kenya) and Kruger National Park. Most stable populations in protected areas in southern Africa; also several populations in eastern Africa. Population declining in eastern, central and western Africa

They are listed by IUCN red list as of Least Concern (2014 assessment) (Bohm and Höner 2015) being widespread in Africa. Decline experienced outside protected areas is not sufficient to warrant listing in a threatened category with global population exceeding 10,000 mature individuals. Within Kenya, spotted hyenas are listed as vulnerable under WCMA, 2013, though scanty information exists on their range or population numbers. They are not CITES not listed.




Figure 15: Map of Hyena range in Kenya

CHEETAH

Acinonyx jubatus



1,160



UPTO 0.9M



UPTO 72KG



EN



DID YOU

KNOW?



Cheetahs must hunt to feed on small to medium sized animals like hares, impalas, wildebeest calves, and gazelles and can attain a full speed of 130 km/ hr and **they don't scavenge.**

3.1.13 Cheetah

i. Introduction



Global population size has been ‘guesstimated’ at 14,000 (Myers, 1975) and ‘less than 15,000’ (Marker, 2002). The species is listed as vulnerable according to IUCN red list criteria (IUCN, 2006a). The most recent estimate by the International Union for Conservation of Nature puts the figure at 6,600 – mainly in eastern and southern Africa – amid fears that the fastest land mammal is racing to extinction. Cheetahs are now extinct in 20 countries and occupy only 17% of their historic range. The remaining populations that are of global importance are found in southern Africa Botswana, Namibia and South Africa and in East Africa Kenya and Tanzania. Scanty information is available about cheetahs in Kenya.

Important populations of cheetahs occur in Tsavo, Mara-Serengeti and Laikipia-Samburu ecosystems. However

the species’ status in other areas is not well understood. Although both cheetahs are economically important to Kenya’s tourism industry, the majority of these animals reside outside the protected areas which are the focus of most tourism. Over 80% of cheetah geographic range, occur on community and private lands. Populations in the protected areas would not be viable if isolated from those outside the protected areas. As such, conservation activities outside protected areas is critical for the long-term survival of the species. Some cheetah populations are transboundary especially along Mara- Serengeti and Tsavo-Mkomazi conservation areas. Transboundary management is key to secure the future of the species in the long term



Figure 16: Map of known cheetah populations

WILD DOGS

Lycaon pictus



DID YOU KNOW?



Wild dogs have **very strong social bonds**, stronger than those of sympatric lions and hyenas thus live in permanent packs consisting of two to 27 adults.



UPTO 1.1M



UPTO 36KG



EN

(Source: <https://drewsproule.com/>)

3.1.14 Wild dog

i. Introduction



African wild dogs are highly social members of the canid family. Packs cooperate to hunt their prey (Creel & Creel, 1995), which consists mainly of medium-sized ungulates. Packs also cooperate to breed, with usually only one female and one male being parents of the pups (Girman et al., 1997a), but all pack members contributing to pup care (Malcolm & Marten, 1982).

Unlike most carnivore species (other than cheetahs), wild dogs tend to avoid areas of high prey density (Mills & Gorman, 1997), apparently because larger carnivores prefer such areas (Creel & Creel, 1996). Lions and hyaenas

are the main causes of death for adult and juvenile wild dogs (Woodroffe et al., 2007a). Probably because of this tendency to avoid larger predators, wild dogs live at low population densities and range widely. Population densities average around 2 adults and yearlings per 100km² (Fuller et al., 1992a) and home ranges average 600-800km² per pack in eastern Africa (Woodroffe & Ginsberg, 1998), with some packs ranging over areas in excess of 2,000km² (Fuller et al., 1992a). Wild dogs occupy home ranges larger than would be predicted on the basis of their energy needs.




Figure 17: Map of wild dog range in Kenya

WATERFOWLS

(FLAMINGOES)

Phoenicopterus roseus



255,936



UPTO 1.5M



UPTO 4KG



LC

DID YOU KNOW?



Flamingos pink colour is obtained from its diet.

Each flamingos' shade of pink varies by species,

with the Caribbean species a vivid pink and the Greater species a pale pink

3.1.15 Waterfowls

i. Introduction



Kenya has one of the most rich avian species diversity in the world with approximately 1,100 species recorded. Out of these, about 170 are Palearctic migrants, moving from Eastern Europe, Russia and Middle East to winter in Kenya, while about 60 are migrants within the African continent. Some 23 are forest dependent species, preferring pristine undisturbed forest habitats. Such species diversity and distribution can be attributed to a diverse habitat type resulting from topographical and climatic variation, resulting from high mountains, the scenic Rift Valley, flat arid and semi-arid landscapes to humid coastal ecosystems. The Kenya's Important Bird Area (IBA) programme has identified 67 important for bird conservation (Birdlife 2021; Figure 34). Most of these sites have also been identified as Key Biodiversity Areas (KBAs), because they are important for the conservation of other biodiversity besides birds.

To enhance our understanding and knowledge on bird distribution in Kenya, The Kenya Bird Map (KBM), a citizen science tool based on reporting of bird sightings across the country has helped map out distribution of birds using bird watchers, including both the local citizen birders and visiting foreign birders at different sites in the country. This method has shed light on bird occurrence and distribution in Kenya from 2013 to date. However it may not bring population status, especially rare birds. The IBA monitoring programme is carried out by individuals, institutions, Site Support groups at IBA sites helps to better understand the conservation status of species and their habitats. However, Bi-annual water bird census has been a key tool in understanding population trends and conservation status of water birds by looking at species and their numbers over the past 30 years. Such information has helped in designating such sites as IBAs by Birdlife International,

Wetlands of international importance (RAMSAR site) as well as UNESCO World Heritage Sites. Information from the biannual waterbird population estimates is used to prepare monitoring reports for the internationally recognised sites. For example, besides being IBAs, Lake Bogoria, Nakuru, and Elementaita are World Heritage Sites using the data, reports and publications from the annual water bird census.

Water birds are key indicators of wetland ecosystem health. They show fast response to changes in wetland habitat quality. Wetlands provide both feeding and breeding sites for both congregating and solitary water birds. Kenya has a rich and diverse wetland habitat ranging from saline lakes within water lakes within the Rift Valley, to fresh water lakes within Lake Victoria forming the larger area. The marine ecosystems along the coastal shore line provide important flyways for migratory birds. The annual water bird census takes place in more than 150 countries around the world, with the sole purpose of monitoring water bird species and population for sound conservation and sustainable management of wetlands. The counts have been on for more than 50 years under the umbrella of the International Water bird Census (IWC) Water bird census in Kenya dates back 30 years of continuous data collection. The counts have taken place on Rift Valley lakes (Lake Bogoria, Lake Nakuru, Lake Naivasha, Olbollosat and Lake Bogoria), the north coastal marine wetlands and constructed wetlands in both Nairobi and central Kenya in Limuru and Thika. This report covers the avian fauna that is found in Kenya Rift Valley lakes of Magadi, Naivasha, Olbollosat, Elmentaita, Nakuru and Bogoria. The water bodies are divided into Freshwater (Lake Naivasha, Nairobi Park, Manguo Swamp,




Figure 18: Important Bird Area sites in Kenya (Source: IBA 2019)





Figure 19: A map showing the distribution of birds' sightings in Kenya using citizen science (Source: Kenya Bird Map, July 2021)

3.2 | RESULTS BY ECOSYSTEMS

3.2.1 Tsavo Ecosystem



In this section we provide results of the Tsavo ecosystem during the 2021 total aerial census. The ecosystem covers an area of approximately 49,611 km². It is the largest continuously savannah range in Kenya representing approximately 38% of the known elephant range in the country. The ecosystem occupies parts of Taita Taveta County, Kilifi County, Kwale County, Makueni County and Kitui County. It borders Kitui County to the north, Tana River County to the east, Kilifi County to the south east, Kwale County to the South, Kajiado County to the west and Makueni County to the northwest. It also borders Tanzania's Mkomazi National Park to the South East forming a key trans-boundary conservation area in Africa (Figure 40).

Previous censuses in the ecosystem indicate it hosts more than one third of the national elephant population (Chase et al., 2016; Ngene, et al., 2017). The ecosystem therefore significantly contributes to the continental range of the African elephant (Thouless et al., 2016). Updates of wildlife populations in the ecosystem have been done over the years using total aerial census (Thouless et al., 2016).

Total aerial census have been implemented in the Tsavo ecosystem since 1962. While not always regularly implemented, the censuses provides a sound long term trend of elephant and other wildlife populations, which has been used for their conservation and management (Ngene et al., 2017). It is important to also mention that the number of wildlife species counted during the censuses have been changing, where in some cases only two (elephant and buffalo), three species (elephant, buffalo and giraffe) and all species larger than dik dik have been counted (Omondi et al., 2002; Kyale et al., 2014; Ngene et al., 2017). The higher number of species counted has been possible because of rigorous training including the use of voice recorders and taking of aerial photos. During the census, data quality was ensured by having a compulsory training of observation, recording and transect test flight for the flight crew. This year's census is the 19th such census in the ecosystem. A good description of the history of the Tsavo aerial census, the methods used over time and the summarized results are outlined by Kyale et al. (2014), Ngene et al., (2013) and Ngene et al., (2017).





Figure 20: A map showing the aerial census area during the Tsavo Ecosystem census, June 2021

3.2.2 Laikipia-Samburu-Marsabit-Meru Ecosystem



Kenya Wildlife Service undertakes wildlife censuses in various ecosystems across the country every 3 to 5 years. The focus for such routine censuses have been within traditional wildlife hotspot areas with various methodologies being employed depending on the target taxa. Aerial counts have been conducted in Kenya since 1960's to count large mammals across selected ecosystems using standard methods (Douglas-Hamilton, 1996, Kangwana, 1996). The Laikipia-Samburu-Marsabit-Meru Ecosystem is home to second largest population of elephants in Kenya (Omondi et al., 2002). The ecosystem generally referred to as Laikipia-Samburu extends far north to Marsabit and South East as far as Meru national park.

Unlike other ecosystems in Kenya, the Laikipia-Samburu ecosystem comprises of predominantly non-formally protected land units; with the community and private land ownership accounting for 97% of the elephant range, making it a human dominated landscape (Ihwagi et al., 2015). In these landscapes dominated by humans, the home range of elephants comprises distinct home ranges connected by migratory corridors through areas (Douglas-Hamilton et al., 2005, Ngene et al., 2010). Loss of habitat through infrastructural developments and change of land use is the most significant threats to elephants in the long term (Nellemann et al., 2013). The increasing human foot print i.e., settlement of nomadic pastoral communities along known historical elephant routes is a major concern. Efforts involving the government, communities and private researchers are under way towards a joint spatial planning that would secure the elephant connectivity. Elephant are the umbrella species and protecting its connectivity routes caters for the needs of other species of top concern in the Laikipia-Samburu ecosystem (Didier et al., 2009). The

Laikipia – Samburu population is subject to a long-term GPS (Douglas-Hamilton, 1971, Thouless and Dyer, 1992, Douglas-Hamilton, 1998, Douglas-Hamilton et al., 2005). Despite the increasing human settlement, elephants are still able to roam freely within Marsabit, Isiolo, Laikipia, Meru and Samburu Counties (Thouless, 1998, Didier et al., 2009, Ngene et al., 2010). However, due to increased human presence often perceived as a risk by elephants, their movement behavior has been shown to shift to moving more at night than during the day when they move into high risk areas (Ihwagi et al., 2018).

The population of elephants in the ecosystem is routinely monitored through periodic total aerial counts (Omondi et al., 2002, Litoroh et al., 2010, Ngene et al., 2013, Ngene et al., 2017). The total aerial elephant census in Kenya are undertaken every three-four year subject to availability of resources, and the past total counts were conducted in 2002, 2008, 2012, and 2017. The area of coverage in 2008 was 46,391km², 2012 was 55,000km² and in 2017 it was extended to cover Meru and Marsabit areas; a total 65,516 km². In addition to these counts covering the entire landscape, a partial-coverage total count focusing on community areas was conducted in 2005 by African Wildlife Foundation. The June 2021 census was conducted as part of the national wildlife census.

Land tenure in Laikipia-Samburu ecosystem is predominantly privately and communal, with only a few land units under formal protection as forest and national reserves (Ihwagi et al., 2015). Land ownership has a significant effect on the potential use of an area for wildlife conservation (Newmark and Hough, 2000, Fitzsimons and Wescott, 2007, Petrzalka and Marquart-Pyatt, 2011), while land use also typically influences the distribution and


abundance of herbivores (Blom et al., 2005, Georgiadis et al., 2007, Ogutu et al., 2009, Ogutu et al., 2014). In turn, animal distribution and abundance can determine the location and intensity of illegal hunting activities (Waltert et al., 2009, Maingi et al., 2012). Land under an official conservation status is traditionally associated with higher protection and abundance of wildlife and is recognized as critical for the conservation of species (Hedges et al., 2005, Pia et al., 2013). In the Laikipia Samburu ecosystem, the Samburu, Buffalo Springs National Reserves and the private sanctuaries in Laikipia on account of their higher protection status host the majority of wildlife population (Ihwagi et al., 2015).

The Laikipia-Samburu ecosystem is home to 93% of global population of Grevy's zebra (*Equus grevyi*), a population that had experienced 85% decline over the last between 1990 and 2017 years (Nelson, 2003; Rowen & Ginsberg, 1992). The last comprehensive census of Grevy's zebra in Kenya was undertaken in the year 2000 (Nelson & Williams 2003), resulting in an estimated national population of about 2,571. In 2004, information contributed by various Grevy's zebra conservation stakeholders suggested that the population had dropped to between 1,567 and 1,976

(Williams & Low 2004). A second stakeholder workshop in 2007 suggested that the number of Grevy's zebra in Kenya was between 1,838 and 2,319 (Mwasi & Mwangi, 2007). However, both the 2004 and 2007 estimates were guesses based on summing tallies over different Grevy's zebra populations. The census conducted in June 2001 incorporated the results from the innovative Gravy Zebra Rally; citizen science entailing taking pictures of all encountered animals and processing the photos through a custom photo recognition database (Rubenstein, pers comms).

The census is designed to sustain the long term aerial monitoring of large mammals while at the same time enrich the species status database. The specific objectives for the aerial census were to: Determine the present status of population of elephants, Determine changes in the elephant population and their distribution since the last aerial census of 2017, Determine population abundance, distribution and trends of other large wildlife within the landscape, Map out various activities in order to assess current pressures on wildlife conservation and provide baseline wildlife data for the country.





Legend

- Flight paths
- Town
- River
- Survey block
- National Park
- National Reserve
- National Sanctuary
- County Boundary



0 25 50
Kilometres

Data Source: KWS/WRTI
Prepared by: F. Muchiri & R. Njeri

Figure 21: LSMM Ecosystem aerial census census blocks, June 2021

3.2.3 Maasai Mara National Reserve (MMNR)- Ecosystem



37,281 ■ Wildebeest	32,358 ■ Zebra Common	11,604 ■ Buffalo
10,610 ■ Impala	8,278 ■ Thomson's gazelle	6,923 ■ Topi
3,893 ■ Grant's gazelle	2,595 ■ Elephant	2,109 ■ Giraffe
1,280 ■ Eland	764 ■ Hartbeest	574 ■ Warthog
534 ■ Hippo	215 ■ Ostrich	162 ■ Baboon
161 ■ Water buck	30 ■ Lesser Kudu	

The 2021 aerial total count of elephant and other large wildlife including the buffalo, giraffe amongst others within the Maasai Mara Ecosystem (MME) was conducted between 6th and 10th May 2021. This report therefore gives a status update of the large mammals populations in relation to human activities within the Kenyan side of the MME. The last status report was produced in 2017, and it documented an estimate of 2,493 elephants, 9,466 buffaloes and 2,607 giraffes in the Mara ecosystem (Mwiu et al., 2017).

The Mara ecosystem being part of the 30,953Km² greater Serengeti - Mara ecosystem is one of Kenya's rich wildlife resource areas and an important agricultural production area. The ecosystem is a major tourism destination due to the diverse wildlife which includes the great wildebeest migration and charismatic wildlife. The wild animals in the Mara ecosystem utilize the PA as well as the surrounding conservancies and dispersal areas (Otichillo et al., 2000).

The ecosystem falls under Narok County which borders Kisii, Nyamira, and Bomet counties to the west, Nakuru

County to the north and Kajiado County to the east. The study area borders Serengeti National Park (SNP) in the Republic of Tanzania to the south, Nyakweri forest to the western region, community owned land coupled with cultivation to the north and loita forest to the eastern regions. The census area is comprised of one PA (Maasai Mara National Reserve (MMNR), community conservancies such as Naboisho, Mara north, Oloisukut, Lemek, Pardamat, Olkinyei, Olarro, Olare orok, Mara Siana, Olderkersi, Enoonkishu and Motorogi which lie in the former community group ranches, and Olchoro oiruha private conservancy among other adjacent community owned land and group ranches. These conservation areas outside the PA include several community and private conservancies demarcated from the former larger group ranches and current existing group ranches that are at various intermediate and advanced stages of land subdivision and human habitation. A detailed review of the historic perspective of aerial census in the ecosystem and a description of the study area is provided by Mwiu et al. (2017).





Figure 22: Map showing the location of Maasai Mara ecosystem

3.2.4 Amboseli-Magadi Ecosystem



28046 Common Zebra	21357 Grants gazelle	10257 Wildebeest	6425 Giraffe
4274 Eland	3358 Impala	2108 Thomson gazelle	1641 Elephant
781 Ostrich	537 Buffalo	344 Lesser kudu	266 Gerenuk
263 Coke's Hartebeest	207 Oryx	165 Warthog	97 Hippo
63 Grey-crowned Crane	32 Waterbuck	15 Dik dik	12 Kori bustard
11 Lion	10 Vervet monkey	8 Reedbuck	6 Secretary bird
2 Bushbuck	1 Duiker	1 Hyena	1 Leopard

The Amboseli-Magadi aerial census area falls in the Kenyan portion of the larger Amboseli-West Kilimanjaro-Magadi-Natron (AWKMAN) Cross-border Ecosystem that covers parts of Southern Kenya and Northern Tanzania. The Amboseli-Magadi aerial census area is located between latitudes 35058' and 3806' south and longitude 1024' and 301' east covering an area of 14,143Km². The study area falls in Kajiado County, which borders, Makueni and Taita Taveta Counties to the East, Narok County to the West, Tanzania to the South and Athi-Kapiti plains to the North. The census area was divided into three sub-regions namely, the Amboseli National Park (ANP) which is the protected core zone, the areas adjoining the park that serve as dispersal areas hereby referred to as the Amboseli dispersal area and Magadi region.

Regular monitoring of wildlife populations is an important activity in conservation of biodiversity. Adaptive management and conservation of landscapes require regular censuses of wildlife abundance, their distributions, and their relation to human activities (Lindenmayer & Likens, 2009). Aerial censuses are efficient and rapid means of acquiring data from large areas of conservation interest and have been extensively used in generating vital information for monitoring wildlife populations. These censuses are aimed at deepening the understanding of spatial and temporal variations in distribution and abundance of wildlife populations as well as some selected human activities in the ecosystem.

From the year 2010, Kenya Wildlife Service (KWS) and Tanzania Wildlife Research Institute (TAWIRI) in partnership with other institutions initiated a collaborative partnership of conducting integrated or synchronized aerial censuses in the Amboseli-Kilimanjaro-Magadi-Natron cross-border

landscape. Under this arrangement, regular wet season (March to May) and dry season (September to October) censuses are conducted with a frequency of every three years (KWS & TAWIRI 2010; Kenana et al. 2010; Kenana et al. 2013) using similar methods. Five censuses have so far been done under this arrangement, constituting two dry and three wet count datasets. To bridge the methodology differences that existed between the two countries, a common methodology has been adopted.

The Kenya Government through KWS and Wildlife Research and Training Institute (WRTI) initiated a National wildlife census in the month of April 2021. This census was therefore done on the Kenyan side of the greater cross border ecosystem which comprised of Amboseli and Magadi census regions covering an approximate area of 14,143 km² (Figure 99)

Ecologically Amboseli-Magadi ecosystem is linked to a cross-border ecosystem in the Tanzania side as described in KWS & TAWIRI (2010). Wildlife population in the ecosystem has been well studied over the past 40 years (Western 1973; Western & Van Praet, 1973; Lindsay 1994; Kikoti, 2009). Most of these studies have, however, focused on smaller portions of the linked ecosystems, mainly confined within national administrative boundaries in Kenya and Tanzania. Consequently, the wider picture of the cross border ecosystem of Amboseli-West Kilimanjaro and Magadi – Natron in terms of wildlife population monitoring had not been adequately investigated until the year 2010. With increasing knowledge of animal movements within the ecosystem and observations of fluctuating animal numbers within the various constituent areas, it is necessary therefore to consistently carry out the broader census in both wet and dry seasons every

Amboseli-Magadi Ecosystem

three years in order to study the changes within the whole area as a unit. Furthermore, adaptive management and conservation of natural ecosystems require effective monitoring of biodiversity, including regular censuses of wildlife abundance and distributions (Lindenmayer & Likens, 2009).

The overall goal for the census was to collect comprehensive wet season data on various wildlife species and human activities within the Amboseli- Magadi Ecosystem (AME). The specific objectives of the census were to:

- Determine wildlife population's abundance and distribution in the Amboseli –Magadi ecosystem
- Determine wildlife population trends in the ecosystem over time
- Determine the extent and spread of human activities in the ecosystem
- Identify threats to wildlife conservation in the ecosystem
- Suggest strategies for effective wildlife management across the ecosystem
- Provide baseline wildlife data for the National Wildlife census





Figure 23: A map of the Amboseli-Magadi Ecosystem

3.2.5 Athi-Kapiti Ecosystem



The Athi-Kapiti ecosystem (AKE) borders Nairobi National Park to the north and constitutes the larger Kitengela-Athi-Kaputiei ecosystem which is bound by Nairobi city in the north, the Kajiado ranches & farms to the west and the Machakos highlands in the north-east and east. It consists of large group ranches, wildlife conservancies, privately-owned ranches and GoK-owned ranches. The ecosystem covers a total of 6510km² and borders Amboseli-Magadi census blocks to the south and Mara census blocks to the northwest.

Even though ground censuses have been conducted in AKE in the past covering a small fraction of the ecosystem, the concept of aerial censuss only took traction as late as last year (2020) where likewise, a small fraction of the ecosystem was covered. By their very nature, aerial censuss are fast, consistent, cost-effective and accurate thereby being the most appropriate method for generating the much needed information including wildlife threats and challenges within the NNP dispersal area.

The aerial census is important to the extent that spatial distribution of threats affecting wildlife conservation in the region would be established. These threats including anthropogenic impacts like human settlements, crop farming, bomas, livestock and quarry/mining among others that substantively affect graze and browse resources base for wildlife species residing in the area. The distribution


and abundance maps of various species in relation to the land use patterns will guide establishment of potential wildlife dispersal and migratory routes.

This census was aimed at mapping out wildlife populations in relation to developments and infrastructure as the city of Nairobi and its suburbs expand in Machakos and Kajiado counties. The specific objectives of the census were:

- To determine the number and distribution of large animals
- To map out various land use types and human activities to assess current pressures on wildlife conservation
- To document the number and distribution of livestock in relation to large mammals in the ecosystem
- To establish spatial distribution of threats to wildlife due to anthropogenic activities
- To interpret the information obtained and deduce sound management decisions to guide management of wildlife in the ecosystem
- To Provide baseline wildlife data for the National Wildlife Census



3.2.6 Lamu-Lower Garissa Ecosystem



Kenya hosts a wide array of biological resources, including wildlife inside of protected areas and outside living in private and community regions. Wildlife is an important revenue generator through tourism and employment for this country and therefore frequent monitoring to establish its status, population and trends provides important information for managers. Furthermore, wildlife resources are faced with many and mounting challenges including mushrooming human and livestock population, climate change that reduces rainfall and alters food resources available, infrastructural developments and human settlements all of which lead to increased fragmentation and habitat destruction. Therefore regular monitoring of wildlife is important.

The most recent census was undertaken in the area in 2015 and 26 mammalian species were counted. The Cape buffalo (*Syncerus caffer*) was the most abundant at 13,754 individuals, followed by the topi (*Damaliscus korrigum*) at 9,686 and reticulated giraffes at 3,254 individuals. The common zebra, warthogs, and common waterbuck were amongst other species counted in this region. The persistence of wildlife in this region is primarily due to the pastoral nature of the communities that leaves wide swaths of land open for grazing. While such practices have proved to be pro conservation, changing lifestyles has led to sedentary settlements of once pastoral communities threatening wildlife conservation. This year's aerial total count was part of the National Wildlife Census (NWS) that sought to enumerate wildlife numbers in all key ecosystems in Kenya to ascertain their status and distribution against the changes in the ecosystems due to human factors.

This is the second aerial total count to be undertaken in the Lamu-Lower Garissa ecosystem after the first aerial census in 2015. Like other ecosystems in Kenya, the

ecosystem hosts a high diversity of wildlife including the buffaloes, topis, giraffes, the critically endangered hirola. Also, it is an important elephant range and past censuses have reported elephant sightings. Wildlife in this region is faced with multiple challenges including insecurity that predispose wildlife to poaching, bush meat, and human-wildlife conflicts. This year's census region included the almost the entire Lamu County and the lower regions of both Garissa and Tana River Counties totaling 23,377km². The team constituted of six aircrafts (2 - cessnas, 1-supercab, and 3-huskys) and lasted eight days. Due to strong winds prevailing in the region, 2km transects were only undertaken during the morning part of the days.

The objectives of the census were to:

- Document the abundance and distribution of all large mammals including elephants, buffaloes, giraffes, etc
- Understand the trend in wildlife numbers in the region,
- Map human activities including fences, logging, settlements, farming, and charcoal burning,
- Provide baseline wildlife data for the National Wildlife census
- Document the distribution and numbers of livestock (e.g. cattle, goats and sheep) and,
- Interpret the information obtained to guide management of elephants and other

Lamu-Lower Garissa and Tana River Ecosystem





Figure 25: Map showing the Lamu-Lower Garissa ecosystem

3.2.7 Naivasha-Nakuru Ranches



The Nakuru-Naivasha Ranches is located in Central Rift valley region and falls within two counties: Nakuru and Kajiado. More than 90% of the conservancies/ranches are found in Nakuru County. The census area is comprised of two Protected areas namely Hell's Gate and Mt. Longonot National Parks. The conservancies in Nakuru County included Eburu forest, Soysambu, Kigio, Malu, Marula Farm, Aberdare Golf Club, Morendat Farm, Karlo lower and upper farms, Kikopey area, Eburu forest, Green Park, Great Rift Vally lodge, Colour Crops, Loldia farm, Muhu Farm, Bilashaka Farm, Rocco farm, Ol-Suswa, Wileli, Korongo farm, Aquilla, Mundui Farm, Oserengoni Conservancy, Hippo point, Crater Lake, Kongoni Game Valley, Olerai, Kedong, Lendolia, Crescent Island, KWSTI, Annex, Game Farm, Sanctuary Farm, Karai area, Lakes Naivasha and Elementaita conservancies including their adjacent riparian areas and flower farms. Among the conservancies covered, only Mt. Suswa Conservancy falls in Kajiado County. The study area lies between longitude 36° 06' 32" E and 36° 36' 33" E and latitude 0° 22' 47" S and 0° 10' 06' 26" S (Figure 138).

The Nakuru-Naivasha ecosystem is rich in wildlife resources. The ecosystem is a major tourism destination due to the diverse wildlife, geo-morphological features, floriculture, geology and other recreational activities. This calls for close and regular monitoring of the wildlife status and distribution. Aerial censuses have not been very common in the area. Alternatively, the area has employed ground animal census since 1999 although hippo census using boats have also been carried out in the area. Most of these ground wildlife censuses have been carried out twice in a year taking into consideration the dry (August-October) and wet seasons (April-July). Bi-annual (January and July) waterfowl counts in the area have also been carried out in the area although these have been extended to cover all the Rift Valley Lakes.

Aerial census was undertaken within the Nakuru-Naivasha ranches in June 2021. The census involved total count for wildlife including the buffalo, giraffe within the Nakuru-Naivasha Ecosystem. We report on the status and distribution on the species of wildlife counted in the ecosystem. This is the first aerial census report since the area has employed ground mammal counts. The census further complements efforts by KWS and partners to attain a comprehensive status of large wildlife populations within the Nakuru-Naivasha area including the aquatic animals such as hippos and crocodiles found in the lakes and rivers. The census aimed to mapping out wildlife populations in relation to developments and infrastructure as the Nakuru-Naivasha Ranches. The specific objectives of the census were:

- To determine the number and distribution of large animals
- To map out various land use types and human activities to assess current pressures on wildlife conservation
- To document the number and distribution of livestock in relation to large mammals in the ecosystem
- To establish spatial distribution of threats to wildlife due to anthropogenic activities
- To interpret the information obtained and deduce sound management decisions to guide management of wildlife in the ecosystem
- Provide baseline wildlife data for the National Wildlife census




Figure 26: Map of the study area showing the protected areas and Conservancies

3.2.7 East Isiolo County



3,000

■ Grants gazelle

788

■ Ostrich

160

■ Reticulated giraffe

81

■ Gerenuk

65

■ Oryx

47

■ Baboon

32

■ Warthog

11

■ Fox

9

■ Dik dik

8

■ Kori bustard

5

■ Lesser kudu

3

■ Impala

2

■ Grevy's zebra

2

■ Common zebra

1

■ Jackal

1

■ Hyena

Isiolo County is one of the counties in the lower eastern region of Kenya. It borders Marsabit County to the North, Samburu and Laikipia Counties to the West, Garissa County to the South East, Wajir County to the North East, Tana River and Kitui Counties to the south and Meru and Tharaka Nithi Counties to the South West. Most of the land in the county is flat low lying plain resulting from weathering and sedimentation. The plains rise gradually from an altitude of about 200 m above sea level at Lorian swamp (Habaswein) in the northern part of the county to about 300 m above sea level at Merti Plateau. The main wildlife species found in the county includes: giraffe, African wild dog, elephant, ostrich, baboons, gerenuks, impala, grants gazelle, leopard, waterbuck, lesser kudu, greater kudu, Grevy zebra, lion just to mention a few. There are over 300 species of birds. The focus of this report is based on census blocks concentrated in Garbatula and Merti sub counties.

Kenya Wildlife Service has been undertaking wildlife census every 3 to 5 years. The focus for such routine census has been traditional wildlife hotspot areas with various methodologies being employed depending on the target taxa. In majority of these ecosystems, aerial census has focused on large mammals namely Elephants, Buffalo, Giraffe, plains and Grevy's Zebra, with different methodologies being employed for other species of conservation concern.

The 2021 National Wildlife Census largely ensured the conventional wildlife census areas are covered and it in addition expanded the scope to include more area and all the species that are observable from air. The census has also in addition included all the human activities within

the landscape. This census is therefore considered the most comprehensive and is expected to guide in holistic understanding of the landscape and wildlife resources therein. Such data is expected to guide on management of wildlife, their habitat and advice on distribution of resources and management strategies. The census methodology followed standard methods described in details by Norton-Griffiths (1978) and Douglas-Hamilton (1996).

The census was proceeded by training to ensure that all pilots and observers understand what is expected of them and that data generated is standardized. Procedures for data cleaning and analysis were followed as outlined in details by Douglas-Hamilton (1996) and Norton-Griffiths (1978). The data was analysed and displayed in maps using ArcGis as outlined by ESRI (2010). Analysis and interpretation of the data was undertaken following procedures described by (1996). The census involved a total of 3 aircrafts. During this time, the landscape was generally dry in majority of the areas and isolated incidences of drought had been reported.

The overall goal for the census was to collect comprehensive data on various wildlife species and human activities within the larger Isiolo sub county. The specific objectives of the census were to: determine wildlife population's abundance and distribution in the census area; establish a baseline for wildlife and human activity dataset for the census area; determine the extent and spread of human activities in the census area; identify threats to wildlife conservation; and, generate data that can assist wildlife management across the landscape.





Figure 27: Location of the study area and Isiolo Census blocks

3.2.9 Lake Nakuru National Park



Lake Nakuru National Park (LNNP) was established in 1961 and is world famous ornithological spectacle and conservation area. Initially, it only encompassed Lake Nakuru and the adjacent 5 km² riparian area adjacent to the lake, but was extended to include a large part of the surrounding savannahs to its current area of 188km². Lake Nakuru is one of the Rift Valley soda lakes at an elevation of 1,760 meters above sea level (m.a.s.l). It lies between Bahati Escarpment on the east and the Mau Escarpment on the west (Montcoudiol, et al., 2019). LNNP has a perimeter fence to contain wildlife as it is surrounded by town centers and communities all round. Fencing has made the park to be a closed island ecosystem (Shah, 2016).

Until recently, the lake had an abundance of algae which attracted a vast number of flamingos that famously lined the shore. It has a watershed spanning over 1800 km² which includes gazetted forests such as Mau, Eburru and Dundori. The Mau Forest consists of plantations and indigenous trees and is part of the Mau complex that covers an area of 650 km². Eburu Forest consists of indigenous tree species covering an area of 87.36 km² to the south of the lake while Dundori Forest in the eastern covers an area of 69.56 km (Shah, 2016).

The park supports a wide ecological diversity with flamingos and other water birds being the major attractions. The ecosystem supports over 300 plant species; 50 different species of mammals; and a variety of terrestrial birds numbering more than 450 species (Hongo, 2020). The park vegetation cover comprises areas of marshland and grasslands with bushy woodland in the rocky cliffs. The park has the euphorbia forest dominated by Euphorbia candelabrum on the eastern side and acacia woodland immediately around the lake shore which is dominated by

Acacia xanthophlea.

The park is a closed ecosystem thus regular wildlife monitoring and counts are important to detect changes in populations and understand the drivers to inform management interventions. Among the challenges that the park faces include: 1) Loss of terrestrial habitat to rising water levels, 2) proliferation of invasive plant species; 3) Overstocking of major herbivores; 4) Susceptibility to wildlife diseases and. 5) Seasonal variations in weather conditions.

For these reasons routine mammal census in the park has been conducted over the years. These censuses have been conducted through road count method which have a limitation in accessing certain parts of the park where the road network is very sparse or in vegetated areas. For the purpose of getting a more reliable estimate of parks buffalo population, a helicopter aerial census was conducted in January 2020 to guide the management in making decision on managing buffalo numbers in the park. Wildlife population assessments are important to enable the national government to understand population dynamics and to assess the success of conservation programs. Data and information generated during censuses assists in making informed decision and ensuring continued attention from the global conservation community. In an effort to understand the status of wildlife in the country, Kenya Wildlife Service with funding from the national government is undertaking a nationwide wildlife census from May –July 2021.

This report covers activities and findings of aerial helicopter census conducted in Lake Nakuru National Park on 4th June 2021. The activity aimed at finding out the density of wildlife population, population size, mapping of

Lake Nakuru National Park ecosystem

wildlife distribution as well as factors that would affect the observed distribution as water sources, habitat conditions and effects of human activities e.g. fishing among other factors

The overall objective of this census was to gather comprehensive information on the current wet season data on the number and distribution of large wildlife species in Lake Nakuru National Park. The specific objectives were to:

1. Determine wildlife population abundance, species composition and distribution in the park
2. Determine wildlife population trends in the park
3. Identify threats to wildlife conservation in the park
4. Provide information for effective wildlife management in the park
5. Contribute data to the development of a national wildlife data portal





Figure 28: A map of LNNP used during the June 2021 wildlife census

3.2.10 Nairobi National Park



1641

Impala

988

Buffalo

546

Coke's Hartebeest

147

Maasai Giraffe

108

Thomson Gazelle

95

Wildebeest

91

Grant Gazelle

89

Eland

40

Warthog

22

Waterbuck

3

Bohor Reedbuck

2


Suni

2

Duiker

1

Bushbuck



304

Common Zebra

32


Hippo

19

White Rhino

10

Black Rhino



135

Baboon

8

Sykes Monkey

5

Vervet Monkey



7

Black-backed Jackal

4

Cheetah

3

Leopard

3


Lion

3

Serval cat

2

White-tailed Mongoose



10

Crocodile

1

Leopard Tortoise



425

Helmeted G. fowl

161

Ostrich

29

Yellow N. Spurfowl

11

Grey crowned crane

7

Secretary Bird

1

Kori Bustard

1

Black B. Bustard

Nairobi National Park Ecosystem

Nairobi National Park (NNP) was established in 1946 as the first National Park in the Country and East Africa. The park covers about 117Km². It is located about 10km from Nairobi City's central business district, the capital city of Kenya. The park is the protected core zone for the wider Athi-Kapiti ecosystem, which is approximately 2000km².

Regular monitoring of wildlife populations is an important activity in conservation of biodiversity. Adaptive management and conservation of landscapes require regular censuses of wildlife abundance, their distributions, and their relation to human activities (Lindenmayer & Likens, 2009). For all stakeholders to manage their wildlife populations effectively, they need to know how many animals they have (the wildlife population); where these animals are found and when (at what time of year) they are found (WWF, 2000). This information will improve problem-animal management activities, increase the productivity of wildlife based enterprises (e.g. tourism/ecotourism) and contribute to improved management of wildlife habitats. In Nairobi National Park (NNP), monitoring of wildlife of populations once every two months has been ongoing for the last three decades spearheaded by Friends of Nairobi National Park (FONNAP) and NNP management. This year the count coincided with, and became part of the National Wildlife Census.

Because of the park's proximity to Nairobi city, impacts of National infrastructural developments, expanding human populations and associated land sub-divisions into commercial and residential plots, have all diminished land wildlife. These changes have made the wildebeest

migration that had over 30,000 animals in the 1960s to completely collapse due to blockage of their migratory routes. In addition, population of warthog (*Phacochoerus africanus*), waterbuck (*Kobus ellipsiprymnus*), hartebeest (*Alcelaphus buselaphus cokii*), and gazelle populations have declined by about 70%. From the bimonthly ground census, it is estimated that 70-80% of the park's animals roam outside of the protected area boundaries to the southern dispersal areas which are still crucial to most of the migratory animals. Unfortunately, those corridors linking the park to the greater Athi-Kapiti plains have been obstructed by development, urban sprawl and land subdivision and sale, or totally blocked by fences. The wildlife population monitoring is therefore important in understanding negative impacts of these developments.

This census aimed to establish the spatial distribution of various wildlife species in NNP and their population trends over the last decade to establish how the wildlife is faring in the face of expanding anthropogenic developments in Nairobi city and its metropolis. The specific objectives of the census were to determine the wildlife species numbers and distribution; determine key wildlife population trends over the last decade; and propose actions to address challenges facing the wildlife populations in the park. The Park has 15 traditional counting blocks based on the parks road circuits and main habitat types. As such, fifteen (15) census groups were constituted to carry out the wildlife census. Each group consisted of at least 4 persons; a team leader, a driver and 2 spotters and was assigned a single block.




Figure 29: Map Showing study area of Nairobi National Park

3.2.11 Oldonyo Sabuk National Reserve Ecosystem



46

■ Vervet Monkey

30

■ Sykes monkey

25

■ Rock hyrax

23

■ Buffalo

19

■ Dik dik

18

■ Olive Baboon

12

■ Colobus monkey

7

■ Bushbuck

4

■ Common duiker

3

■ Tree Hyrax

Oldonyo Sabuk National Park (OSNP) lies about 80 kilometers east of Nairobi city and 27 Kilometers from Thika Town off Thika-Garissa road. It is situated in Machakos County and lies between latitudes 1005' and 1010'S, longitudes 37010' and 37020'E at 2145M above sea level at the peak of the mountain (Figure 187). The park was gazetted in 1967, covers about 20.7 km² and is entirely a mountain. Four main vegetation types occur in the park with the most dominant being the closed bushland followed by closed trees and the open grassland/bushland in addition to the rain-fed herbaceous cover. The climatic conditions vary over time and space in the area. Other than the volcanic formations of granite rocks, the soils are deeply weathered, except where eroded on steep slopes or where un-weathered rock outcrops occur.

As in all other ecosystems with wildlife, the park is declining in ecological status at an alarming rate as a result of extensive agriculture, industrialization and other anthropogenic activities around it. These activities that influence the park negatively include charcoal burning, tree logging, poaching, quarrying, illegal grazing and frequent arson attacks. All these are adversely affecting wildlife in the park by threatening their survival. There is therefore need for accurate data on all wildlife species especially their population status and trends so as to make recommendations for management actions/decisions.

Regular monitoring of wildlife populations is an important activity in conservation of biodiversity. Adaptive management and conservation of landscapes require regular censuses of wildlife abundance, their distributions and status of their habitats in relation to human activities (Lindenmayer & Likens, 2009). For all stakeholders to manage their wildlife populations effectively, they need to know how many animals they have, where these animals are found and at what time of year they are found (WWF, 2000). This information will improve problem-animal management activities, increase the productivity of wildlife based enterprises (e.g. tourism/ecotourism) and contribute to improved management of wildlife habitats. In Oldonyo Sabuk National Park (OSNP), these regular censuses have largely been lacking owing to constrained resources and difficult terrain. Indeed, the current wildlife census in OSNP serves as the only record about the status of wildlife in the park. The only other record was a Biological Resource Inventory done in November, 2007.

The goal of the census was to establish a baseline of the population status and distribution of various wildlife species in OSNP. This goal was achieved by the following specific objectives: to determine the number and distribution of different wildlife species in the park and identify the status and impact of illegal activities on park's wildlife and habitat.





Figure 30: Map showing the locatipon of Oldonyo Sabuk National Park, Machakos County, Kenya

3.2.12 Mwea National Reserve Ecosystem



Mwea National Reserve (MNR) was gazetted as a National Reserve (NR) by legal notice No. 2 of 29th January 1976. The gazetted area was initially 68km², but later in year 1978 excisions for settlements were done and only 42km² remained as a National Reserve (Figure 191). The reserve is managed and protected by Kenya Wildlife Service (KWS) on behalf of the County Government of Embu (CGE). In 1998, a perimeter electric fence was erected enclosing the reserve areas where it borders the local communities' land. The fence serves a dual purpose of mitigating wildlife conflict by preventing crop damage by animals as well as limiting unauthorized access into the reserve which may lead to destruction of habitat and poaching. MNR is actively managed to achieve desired conservation goals and safeguard the welfare of the neighboring community. To meet the conservation goal, other management practices have been implemented that include managing animal population through re-introductions and translocation.

MNR has been home to elephants and other herbivores for a long time. In the year 1979 it was estimated that there were 49 elephants in the reserve. These elephants were believed to move freely within the reserve, Tana River, Kiambere forest and North Kitui National Reserve (Ngene et al., 2012) (Njumbi, 1995) (Litoroh, Nicholas, & Masinde, 1994). Increased human settlements and agricultural

activity around the reserve, and the subsequent construction of an electric fence, led to isolation and confinement of majority of wildlife. The enclosure also led to various challenges. For example, the occurrence of anthrax outbreak in 2011 that resulted in the death of 11 Nubian giraffes (*Giraffa camelopardalis Nubiani*) and one lesser Kudu (*Ammela phusim berbis*; Kaitho et al, 2013). Local extinction of some wildlife species like Kongoni, common zebra, giraffe and ostrich have been reported which hitherto used to be abundant. These challenges have prompted the re-introduction of certain animals, for example the Nubian giraffe in 2009 (Njumbi et al., 1996) and various zebra introductions. However, despite the numerous challenges, some wildlife species have been thriving. The elephant population for example has continued to grow over the years leading to 23 individuals had to be trans-located to Tsavo East National Park in 2007.

Several aerial and ground wildlife censuses have been undertaken to establish population status of wildlife in MNR. These however have focused mainly on the large mammals (e.g., the recent census of year 2012 and year 2017). The year 2021 aerial census attempted to establish the population status of a broader number of species that were observable from air.




Figure 31: Location of Mwea National Reserve and basic infrastructure within the reserve

3.2.13 Ruma National Park



550 ■ Giraffe	365 ■ Buffalo	244 ■ Zebra
56 ■ Defassa Waterbuck	38 ■ Impala	33 ■ Baboon
28 ■ Topi	21 ■ White Rhino	14 ■ Black Rhino
12 ■ Lelwel's Harte Beast	8 ■ Roan Antelope	5 ■ Oribi
3 ■ Bush Buck	3 ■ Bohor Reedbuck	1 ■ Crown Cranes
1 ■ Grey Bush Duiker	1 ■ African Fish Eagle	1 ■ Black Bellied Bustard

Ruma National Park is located in Lambwe Valley between Ndhiwa, Suba North and Suba South Sub-Counties in Homa Bay County, Western Kenya. As at 1966 the gazetted Lambwe Valley Game Reserve had an area of approximately 308 km². The park was gazetted under Legal notice no 100 of 14th June 1983 as a National Park from its previous protected area category as Lambwe Game Reserve, with the rea coverage being 120km².

The Lambwe valley floor which forms Ruma National Park is sandwiched by Gwasi Hills to the west and Kanyamwa Escarpment to the east and other satellite geological features such as Gembe Hills and Ruri Hills to North West and North of the Park respectively. Adjacent lands provide rich culture and history of the neighboring Luo and Abasuba communities with land use mainly being subsistence and commercial crop cultivation and livestock keeping.

Wildlife population censuses and assessments are important management tools enabling national governments to understand population dynamics and evaluate success and failure of conservation programs. Data and information generated from these censuss

assists in making informed decision and ensuring continued attention from the global conservation community. It is because of this that Wildlife Research and Training Institute (WRTI) in collaboration with Kenya Wildlife Service endeavors to understand the status of her wildlife by carrying out nationwide wildlife census.

This report covers activities and findings aerial census conducted in Ruma National Park on 31st May 2021. The overall objective of this census was to gather comprehensive wet season data on the number and distribution of large wildlife species in Ruma National Park. The specific objectives were to:

1. Determine wildlife population abundance, species composition and distribution in the park
2. Determine wildlife population trends in the park
3. Identify threats to wildlife conservation in the park
4. Provide information for effective wildlife management in the park
5. Contribute data to the development of a national wildlife data portal





Figure 32: Map of Ruma National Park showing flight path

3.2.14 Shimba Hills Conservation Areas



The Shimba Hills Conservation Area consists of the Shimba Hills National Reserve, Mkongani (west and north) Forest Reserves, and Mwalunganje (Elephant Sanctuary and Forest Reserve; Figure 201). These conservation areas are home to the remaining population of elephants in Kwale County. Before the elephants were confined to the conservation area, they used to roam freely within the Kwale County (Estes 1974). Prior to 1960s, the elephants were part of a larger population that ranged across the entire Kwale County and as far as Tsavo and Mkomazi, which are about 60km north and 40km south respectively (Estes 1974). Huge herds of the elephants annually migrated every October and November from the south to the east of the conservation areas in search of palm seeds. However, controlled shooting, human settlements and poaching gradually eliminated the elephants from the rest of the county (Thouless et al. 2008). For instance, between 1961 and 1962, about 250 elephants were shot in the area to create the Shimba Settlement Scheme (Estes 1974).

Human population has continued to increase in the area after the establishment of the settlement scheme. This was followed by a wave of poaching in 1970s and 1980s, which forced the remaining elephants to be confined in the conservation areas for protection (Thouless et al. 2008). At the same time, human-elephant conflicts started to be experienced in the area as human population and area under farms increased. In order to mitigate human-elephant conflicts, the conservation areas within the Shimba Hills were fenced with electric fence between 1991 and 1995 (Thouless et al. 2008). Fencing resulted to confinement of elephants within the conservation area. The confinement was a threat to the woody vegetation as elephants at higher densities started to destroy them (Ross 1984). As a result, it became necessary to constantly monitor the elephant population within the Shimba hills conservation areas.

Elephant population monitoring in Shimba hills has followed three main methods, which include aerial

count (Ngene et al., 2013; Tamoooh, 2009; Kimtai, 2007; Kahumbu, 1997; Litoroh, 1997; Kiiru, 1995), dung count (Omondi, et al., 1998;; Mwathe, 1995; Reuling et al., 1992) and individual recognition (Kahumbu, 2002). Ngene et al. (2013) estimated the elephant population in the Shimba hills conservation areas at about 274 animals, and provides a summary of past elephant population estimates in the area. Using the dung count method, Reuling et al. (1992), Mwathe (1995), Omondi et al. (1998) and Litoroh (2002) estimated about 429 ± 128 animals, 453 ± 181 animals, 598 animals, and 649 ± 151 animals respectively in the conservation areas. Kahumbu (1995), Litoroh (1995), and Kimutai (2007) using a helicopter sighted a minimum of about 467 and 305 elephants in the conservation areas. However, Kahumbu (2002) estimated 658 animals using the individual recognition method after removal of 30 bulls from the conservation areas (Muir, 2000). In 1995 and 2009, about 232 and 144 elephants were sighted in the conservation areas using a fixed wing (Kiiru, 1995; Tamoooh, 2009).

The aerial census aimed at updating the population status of elephants and other mammals in the Shimba Hills Conservation Areas. The specific objectives of the aerial census were to: establish the number and distribution of elephant and other mammals in the Shimba Hills conservation areas; establish the number and distribution of elephant carcasses; and, establish the human influence specifically distribution of logging activities in the conservation areas. It is important to continue monitoring elephants and other large mammals' population in the Shimba Hills Conservation Areas. The data and information obtained will illustrate the number and distribution of elephants and other large mammals in the conservation areas. This information is important to conservation area managers and policy makers as it will guide them in making decisions regarding future management of elephants and other large mammals in the conservation areas in Shimba Hills.





Figure 33: Shimba Hills Study Area

3.2.15 Tana River Remnant Forests



The Tana River, which is the longest river in Kenya has its source from Mt. Kenya and the Aberdares draining to the Indian Ocean at Kipini (Figure 203). The river flows through a broad flood plain of alluvial sediments in its lower reaches primarily covered by grass and numerous patches of forests and woodlands. The flood plain forest can be divided into two; the extensive forests north of Garsen and the small fragments occurring along the river courses south of Garsen. The river mouth at Kipini is dominated by the tall mangrove forests represented by the eight mangrove species found in Kenya. The middle and lower sections of the river is dominated by three communities including the Pokomo who are agriculturalists, Orma and Somali who are pastoralists.

The lower Tana River forests are home to several primate species including the endangered Tana River Red colobus (*Piliocolobus rufomitratus*) and the Tana River Mangabey (*Cercocebus galeritus*) (Wahungu et al., 2005). The two species have lost more than 70% of their primary riverine forest habitat because of the anthropogenic activities mainly forest clearing for Agriculture (Moinde-Fockler et al., 2007; Kivai, 2018). According to Moinde-Fockler et al. (2007), over a period of 21 years (1979-2000), forest cover amounting to 29% inside, and 38% outside of the TRPNR has been lost. This loss of habitat has been associated with the sharp decline in numbers and the last detailed census put the average global population of Tana River red colobus at 788 individuals and Tana River mangabeys at 2069 individuals (Karere et al., 2004). Six other nonhuman primates inhabiting the region include the yellow baboon (*Papio cynocephalus ibeanus*), lowland sykes monkeys (*Cercopithecus mitis albotorquatus*), vervet monkeys (*Chlorocebus pygerythrus hilgerti*), garnett's galago (*Otolemur garnetti lasiotis*), Senegal galago (*Galago senegalensis braccatus*) and Kenya coast galago (*Paragalago cocos*) (Suleman, et al., 2001).

The Tana River red colobus and Tana River mangabey are endemic to Kenya, occurring in isolated populations in 27 and 30 patchy and highly fragmented gallery forests respectively. They occur within a geographical short range of about 60 km that supports approximately 80 forest patches along the lower Tana River (Butynski and Mwangi, 1994; Karere et al. 2004). Most of these forest patches however occur north of Garsen-Witu road to TRNPR with a few records 5-15Km south of this road at Onkolde and Masewe forest patches (Hamerlynck et al., 2012; Karere et al., 2004). Increase in community needs for the forest resources within these forest patches has raised the concern of understanding the conservation status of the diverse fauna. Several primate censuses have therefore been undertaken over the last few decades to determine the Tana River mangabey and Tana River colobus distribution and demography (Groves et al., 1974; Andrews et al., 1975; Marsh, 1978; Kinnaird and O'Brien, 1991; Decker and Kinnaird, 1992; Kahumbu and Davies, 1993; Butynski and Mwangi, 1995; Muoria et al., 2003). All these census targeted different forest patches with Marsh 1975 census being the first detailed census where 59 of the known 77 forest fragments was covered (Marsh, 1978). Using similar methods, Butynski & Mwangi (1994) covered 60 of the forest fragments while Karere et al., (2004) covered 73 forest patches in 2001. The 2001 census forms the basis around which the current population trends for the two endangered primates will be assessed.

Despite, the negative impacts on the lower Tana ecosystem, information is lacking on the current population dynamics of the Tana River mangabey, the Tana red colobus and changes on their habitat status. The last meaningful habitat and population assessment of the two primates was conducted in 2001, and unfortunately, due to community resistance and protest against the GEF project, mass destruction of the habitat ensued. Nonetheless, the

Tana River Remnant Forest Ecosystem

impact of such forest changes on the endangered Tana River primates and their habitat is yet to be established. Thus, lack of continuous monitoring and documentation of the effects of forest cover change on Tana primates has contributed to the conservation challenges facing the two species. This census was therefore undertaken to better understand the current population status and distribution of the critically endangered Tana River mangabey and the Tana River Red colobus and strengthen the long-term conservation of the two species and their habitat.

The main purpose of undertaking this census was to understand the current population status of the critically

endangered Tana River red colobus and Tana River mangabey and other wildlife species inhabiting the Tana River flood plains by evaluating the effects of habitat changes on population numbers and distribution

- i. To determine the current population size, distribution and density of the Tana River red colobus and Tana River mangabey, which are sympatric in the lower Tana River forest patches
- ii. To establish the forest habitat changes for the two species over the last 20 years and map out the current habitat suitability for the two species.





Figure 34: Map of the forest patches along the Lower Tana River extending from Nkanjoja to Onkolde indicating the location of TRNPR and including the un-censused section of the middle sector. Adopted from Butynski & Mwangi (1994)

3.2.16 Mandera County



Kenya's rich wildlife resource is one of the key economic pillars of the country. However, the country is not receiving optimized benefits as anticipated due to inadequate knowledge on the status of the country's Wildlife populations as there is no comprehensive scientific data for informed policy direction. To achieve this, there is need for a coordinated National wildlife census in all conservation areas to support implementation of GOK conservation policies and development of evidence-based decision support tools for adaptive management, integrated cross sectoral and multi-scale planning for conservation and sustainable development to optimize wildlife benefits to the country.

The number and distribution of species listed in schedule six (6) of the Wildlife Conservation and Management Act, 2013 (WCMA, 2013) whose numbers and range have significantly declined in the past three decades require regular monitoring. Such wildlife Censuses have become a legal requirement. Every three to five years the Ministry in-charge of wildlife is required to provide species data and information as outlined in the WCMA, 2013 to develop the status of wildlife conservation report as well as the status of wildlife resources monitoring report after every five years. These reports are then presented to parliament by the Cabinet Secretary Responsible as stipulated in sections 49(4) and 64(3) of the WCMA, 2013.

Mandera County wildlife sample count was last done in 2011 by department of resource census and remote

sensing(DRSRS)and since then policy decisions on wildlife conservation in the County has relied on the data. The data obtained from Mandera census is therefore aimed to be manager's tool to manage wildlife resources in the County and manipulate the range condition and productivity of the various wildlife species, including distribution and provision of space and grazing. It will also be useful in provision of security towards wildlife while at the same time guide tourism activities in the conservation areas. Issues of human-wildlife conflicts will also be adequately addressed as well as resource allocation towards conservation and monitoring of the same.

The goal of the sample aerial census was to establish a wildlife population status in Mandera County. The specific objectives of the census were to: determine wildlife population abundance and distribution; assess the wildlife population trends over time; and, identify threats and recommend appropriate strategies to wildlife conservation and management in the County.

The county is located at the extreme end of North-Eastern Kenya, bordering Somalia and Ethiopia and is about 25,991km². It lies between latitude 20 11' North and 40 17' North and longitudes 390 47' East and 410 4.8' East. Administratively, the county is divided into seven (7) sub-counties namely: Mandera East, Mandera North, Mandera west, Lafey, Banisa and Kutulo (Figure 204).





Figure 35: A map of Mandera County showing the constituencies

3.2.17 Garissa County



The census was carried in Garissa county, which lies between latitude 10 58'N and 201'S and longitude 380 34'E and 410 32'E and covers 44,174.1km². It borders the republic of Somalia to the East, Lamu County to the South, Tana River County to the West, Isiolo County to the North West and Wajir to the North (Figure 211).

Garissa County, in the former North Eastern, is one of the rangeland counties found in Kenya. Kenya's rangelands constitute the arid and semi-arid lands (ASALS). The rangelands are dry, hot and have erratic rainfall often below 600mm annually and the home to almost 20% of the country's population mainly the pastoral communities with their large livestock populations. The rangelands are therefore very important in the Kenyan economy in terms of livestock production, as well as the conservation and management of wildlife that occurs in abundance and diversity. It is estimated the ASALS support almost over 50% of the nation's livestock population and over 70% of the country's wildlife population. In most of the year, over 70% of Kenya's terrestrial wildlife reside outside the protected areas and compete with livestock for resources such as water and forage. But in other cases, wildlife is restricted in the parks/reserves thereby degrading their habitats, particularly the vegetation.

Wildlife and livestock in the county is continuously facing increasing risks such as drought, insecurity, animal diseases, increasing human populations and land fragmentation. However, wildlife species co-exist and compete with livestock for survival. To address these challenges, there is need to generate adequate knowledge on the status of the wildlife and livestock populations for informed decision making.

The aerial sample count for Garissa County was part of the National wildlife census in all conservation areas to support implementation of GOK conservation policies and development of evidence-based decision support tools

for adaptive management, integrated cross sectoral and multi-scale planning for conservation and sustainable development to optimize wildlife benefits to the country.

The wildlife Act 2013, schedule six (6) requires production of regular reports on the status of wildlife species whose numbers and range have significantly declined in the past three decades. These reports are presented to parliament by the Cabinet Secretary Responsible as stipulated in sections 49(4) and 64(3) of the WCMA, 2013.

Garissa County wildlife sample count was last done in 2011 by department of resource census and remote sensing(DRSRS) and since then policy decisions on wildlife conservation in the County has relied on the data.

This census is therefore targeted to be a manager's tool to manage wildlife resources in the County and manipulate the range condition and productivity of the various wildlife species, including distribution and provision of space and grazing. It will also be useful in provision of security towards wildlife while at the same time guide tourism activities in the conservation areas. Issues of human-wildlife conflicts will also be adequately addressed as well as resource allocation towards conservation and monitoring of the same.

The goal of this Census is to establish a wildlife population status in Mandera County. The specific objectives that aims to achieve this goal are to:

- Determine wildlife population abundance and distribution
- Assess the wildlife population trends over time
- Identify threats to recommend appropriate strategies to wildlife conservation and management in the County




Figure 36: A map of Garissa County showing the constituencies

3.2.18 Wajir County



2,329,507
■ Shoaat

261,327
■ Camel

192,854
■ Cattle free ranging

14,684
■ Grant Gazelle

12,967
■ Donkey

7,473
■ Gerenuk

6,120
■ Reticulated giraffe

3,090
■ Oryx

3,009
■ Ostrich

2,322
■ Warthog

302
■ Grevy zebra

201
■ Lesser Kudu

141
■ Lion

40
■ Dik dik

Wajir county is located in the North-Eastern region of Kenya between latitudes 10N60'N and 0020'N and latitudes 390E and 410E. The County is a cross border county located in the north eastern region of Kenya. It borders Republic of Ethiopia to the North and Somalia to the East, Mandera County to the North East, Marsabit to the West, Isiolo to the South West and Garissa to the South. The census focused on the entire county which is 55,840km². The County is divided into 6 constituencies as shown below. Wildlife is an important natural resource in arid and semi-arid zones of Kenya and its management requires regular species counts to provide a basis for measuring population changes over time and space.

Wajir County falls within the semi-arid pastoral rangeland of Northern Kenya, where wildlife and livestock abundance are inversely correlated with livestock concentrated around permanent water sources and wildlife density and diversity increasing much further from water points. This inverse correlation between livestock and wildlife abundance has been attributed to indirect interference from associated with livestock most likely because of poaching and harassment, rather than direct competition for forage (Augustine; 2010).

The coat of arms for the county is a reticulated giraffe and a camel and the county is considered as the home and a stronghold of the two species. From 1977 to 2017, the Department of Resource Census and Remote Sensing (DRSRS) has consistently conducted aerial sample counts of livestock and wildlife in the North-Eastern region of Kenya aimed at assessing the status of the resources to inform decision making. According to the National Recovery and Action Plan for giraffes in Kenya (2018-2022), records indicate that, there is a 67% population decrease of giraffe since 1970s. This decline has affected all the three recognized subspecies and addressing

this decline is of high priority. From 13th to 23rd June 2021, Kenya Wildlife Service (KWS), Wildlife Research and Training Institute (WRTI), DRSRS and National Air Support Department (NASD) jointly carried out a wildlife sample Count in Wajir County as part of the National Wildlife Census with the aim at updating the status of wildlife species in the region

Records of reticulated giraffe species indicate that the species has recently suffered a major and rapid decline giving rise to concern about its long term survival. According to the National recovery and action plan for giraffe in Kenya (2018-2022), Wajir County had an estimated population of 3,797 reticulated giraffes in 2016. All the three giraffe species in Kenya are protected under the Wildlife Conservation and Management (WCMA) Act 2013. Schedule 6 of the Act lists Nubian's giraffe as endangered while the other two species (Maasai and Reticulated) are listed as vulnerable and hence the need for action towards conservation of the three species.

Threats facing this species are both intrinsic and extrinsic in nature. The extrinsic threats include: poaching/illegal hunting, habitat loss, climate change, infrastructural developments and inter-specific competitions while intrinsic threats include; inbreeding, dietary complications (e.g., toxins) and diseases.

The giraffe recovery and action plan calls for species censuses in areas where information is lacking or gaps exist to assist in mapping range boundaries and distribution of giraffe subspecies in the country. This census was therefore aimed towards provision of information to guide decision making and consequently save the giraffe population by addressing the threats faced by the species. Increased data collection is also expected to raise awareness on the plight of the species in this region.

Wajir County Ecosystem

The goal of the census was to determine the abundance and distribution of large and medium mammals in Wajir County. The specific objectives were to:

- i. Determine the number and distribution (range) of wildlife species in Wajir County.
- ii. Assess the distribution of available natural resources in relation to wildlife including water and forage.
- iii. Map habitat destructive (logging, charcoal burning) and human activities (farming, settlements) in relation to wildlife distribution and abundance
- iv. Map the distribution and abundance of livestock in relation to large mammals in the ecosystem





Figure 37: A map of Wajir County showing the location of the constituencies

3.2.19 Turkana County



The Kenya rangelands contain substantial numbers of wildlife. Wildlife contributes substantially to the national economy through tourism as the prime tourist attraction. Wildlife based activities contribute approximately 70% of gross tourism earnings. Generally, populations of many wildlife species in the country have declined over the years owing to factors such as habitat loss, changing land use patterns, poaching and drought, including other climate change related factors.

The County shares Lake Turkana with Marsabit County. The total area of the county is 77,000 Km² and lies between Longitudes 34° 30' and 36° 40' East and between Latitudes 10° 30' and 50° 30' North (Figure 229). Turkana County is home to several wildlife species. Some of the species that were observed during the count include: the elephants, gerenuk, lesser kudu, giraffe, Oryx, Grant's gazelle, Duiker, and Bushbuck, among other mammals.

The decrease in numbers for some wildlife species over the years as shown in trend analysis could be attributed to negative impact resulting from climate change, poaching and competition for resources (forage and water) with both livestock and humans, and disruption by increased human activities. Conservation of wildlife in protected areas such as South Turkana National Reserve should continue. Kenya Wildlife Service (KWS) may enhance anti-poaching measures to stem poaching of some species. Also, involvement of local communities in wildlife management can be incorporated in the wildlife conservation strategy of the county.

Livestock grazing is the main land use and economic backbone of Turkana County. Therefore, there is need to support the pastoralists so as to promote their economic base and enhance their livelihoods. This could be achieved

through increased infrastructure development; provision of marketing facilities and construction of abattoirs for livestock off-take during drought to prevent losses; improved livestock breeds; sustainable range utilization during the wet seasons, and provision of insurance for the risks of livestock deaths during adverse weather conditions, among other interventions.

The key threat to wildlife conservation and management in the county is land use / land cover changes. This is especially more so due to increased human related infrastructure development. In recent years, there has been enhanced mining and exploration of oil and other minerals in the county. This has led to increased human population from prospectors and visitors, and also interference with wildlife habitats.

Although the county has two National Reserves in existence, publicity and public awareness about Nasolot and South Turkana National Reserves is minimal and they rarely attract visitors. There is need for integrated management of these two reserves between KWS and the County Government of Turkana.

The goal of this census was to determine population size and distribution of large to medium sized mammals. The specific objectives were to:

1. Determine wildlife population abundance, species composition and distribution in the County
2. Assess the wildlife population trends in the County
3. Identify and map threats to wildlife conservation in the County
4. Map the distribution and abundance of livestock in relation to large mammals in the County





Figure 38: Map showing the location of Turkana

3.2.20 Marine ecosystem



Kenya is endowed with biologically rich marine ecosystems ranging from, mangrove forests, coral reefs, seagrass beds, estuaries, sandy shores, sand dunes and rocky shores. These ecosystems provide important goods and services such as; habitats for fish and other aquatic and terrestrial organisms, erosion control, provision of wood and non-wood forest resources, provision of food, water and industrial resources to millions of people along the coast. This chapter highlights the status of these ecosystems and their associated biodiversity; pressures that threaten their long-term integrity; as well as management interventions.

The information provided covers approximately 600 km long coastline extending from the Kenya-Somalia border at Ishakani in the north (1.7°S; 41.5°E) to Kenya-Tanzanian border at Vanga in the south (4.7°S; 39.2°E; Figure 1). The Kenyan territorial waters extend from the shoreline up to 12 nautical miles covering approximately 9500km². The Exclusive Economic Zone (EEZ) which extends to the 200 nautical miles offshore covers an approximate area of 142,000 km². The coastal counties include: Lamu, Tana River, Kilifi, Kwale and Taita- Taveta.

The Kenyan inshore waters experiences semi-diurnal tides with a spring tidal range of not exceeding 4 m. The coastal offshore waters experiences swell whose magnitude varies

in different periods of the year. The waves are usually very large with a maximum significant height of 8m during the southeast monsoon (May-October) approaching the coast predominantly from south-east and southwest direction.

Kenya hoists a number of marine species of conservation concern due to their rarity, endemism and have evidence of declining population. They include marine mammal species (Cetaceans and Sirenians), sea turtles and bony fishes and elasmobranchs (sharks and rays). A total of 24 marine species have been reported in inshore and off shore waters of Kenya. The most frequently reported species are humpback whale during their peak migration period, July to September and the Indo-Pacific bottlenose dolphin from November-April. Humpback whale sightings are higher because of increased observer effort during the migration period.

The objective of this section is to provide the current information and status of marine biodiversity focusing on marine habitats and species of conservation concern which include Marine mammals, Sharks and Ray, Sea turtles, Coastal birds, Exploited fishery resources, Corals reef ecosystem, Seagrass ecosystem and Mangrove ecosystem.





Figure 39: Map showing the location of Marine

3.2.21 Nasalot-South Turkana-Kerio Valley ecosystem



The Nasalot-South Turkana-Kerio Valley ecosystem covers Nasolot, South Turkana, Rimoi and Kamnarok National Reserves and the surrounding areas with an estimated area of 4571km². It encompasses elephant ranges of West Pokot, Elgeyo Marakwet, Turkana and Baringo Counties. This dispersal area covers a part of the Kenyan North Rift between 204' N and 0046' North and 3503'E and 3602' East. The ecosystem comprises the Kerio valley which is lined by the Turkwel gorge to the North. The gorge branches into escarpments flowing southwards to Keiyo and Baringo which host Rimoi and Kamnarok National Reserves respectively. Nasolot and South Turkana National Reserves covering 92 km² and 1019 km² respectively form the Northern part of the study area while Rimoi and Kamnarok National Reserves in the south cover 66km² and 87.7km² respectively. The Kerio valley is situated at 1,000m and is formed by a narrow and long strip of approximately 80 km and by maximum 10 km wide (OP/MPND, 1991 & 2000; Chebet and Dietz, 2000; Jeatzold and Schmdt, 1983). The total aerial census took place in this area as part of the National Wildlife Census.

Kenya through the Wildlife Research Training Institute (WRTI) and the Kenya Wildlife Service (KWS) is for the first time undertaking a one-off national wildlife census to establish the status of her wildlife resources. Kenya's rich wildlife resource is one of the key economic pillars of the country. However, the country is not receiving optimized benefits as anticipated due to inadequate knowledge on the status of the country's wildlife populations as there is no comprehensive population data for many species of wildlife to inform management.

In every three to five years the Ministry of Tourism and Wildlife is required to provide information as outlined in the Wildlife Conservation and Management Act (WCMA),

2013 as well as the status of wildlife resources monitoring report respectively. These reports are supposed to be presented to Parliament by the Cabinet Secretary responsible for Wildlife Conservation and Management as stipulated in sections 49(4) and 64(3) of the WCMA, 2013. This report focuses on a two day aerial census which was conducted in Nasolot/South Turkana/Rimoi/Kamnarok areas on 4th and 5th June 2021. South Turkana and Nasolot reserves probably hold the largest elephant population in western Kenya.

This census will enhance conservation efforts by determining the exact numbers of all wildlife including the endangered species and their exact location. It will also establish an updated data base which will form a basis for establishing real economic value of wildlife capital for reflections in the national budgeting process and enhance conservation efforts.

The last comprehensive total aerial count was carried out in 2015 (Chase et al., 2015). In 1973 it was estimated that there were 1,500 elephants in Turkana District, but figures were unavailable for West Pokot, Elgeyo Marakwet and Baringo districts (Jarman, 1973). Censuses undertaken by Department of Remote Sensing and Resource Censuss (DRSRS) in 1970's and 1980's generally yielded low estimates apart from one count in 1981 that recorded more than 800 elephants outside the reserves. Kenya Wildlife Service estimated in 1990 that 400 elephants utilized Nasalot and south Turkana with another 100 in Rimoi and Kamnarok. DRSRS sample counts in the same year gave estimates of 535 for West Pokot (including Nasolot), zero for Turkana and 596 for Baringo. In a sample count that was undertaken by Mbugua (1992) in Elgeyo Marakwet, a small number of elephants was sighted inside the sample strips, and thus the count was treated as a

low-intensity total count. The count further presented a similar figure of 580 elephants, of which 525 were in Nasolot and South Turkana. Since this was carried out at a low intensity, it was estimated that the total population could have consisted over 900 elephants. Total aerial counts of the Nasolot/South Turkana elephants were carried out in June 1997 (Muriuki et al.1997) and July 1999 (Omondi et al.1999) that employed methods by Douglas-Hamilton et al.1994 and Douglas-Hamilton, 1997. A total of 852 and 792 elephants were counted in 1997 and 1999 respectively. In both counts the majority of elephants were found in the northern block of Nasolot/South Turkana, with smaller numbers observed in the southern Kerio block in the environs of the Rimoi and Kamnarok National Reserves. In 2002 (Omondi et al, 2002), a total aerial count was conducted and a total of 490 elephants were recorded. Most of them were found further south east towards South Turkana National Reserve and the rest in the furthest south of the Kerio valley in Rimoi National Reserve. The most recent total aerial count was undertaken in 2010 (Edebe et al, 2010) and a total of 362 Elephants were recorded. Other wildlife species counted included grant gazelle, lesser kudu, dik dik, and domestic livestock mainly cattle, camel, sheep and goats (Shoats). Cultivation and new human settlements were observed while pastoralism was noted as the main economic activity.

The objective of the exercise was to ascertain Nasolot-South Turkana-Kerio valley wildlife population and distribution, determine the exact locations of wildlife in those areas to minimize human wildlife conflict and identify threats to wildlife conservation and management.

The specific objectives of the census were:

- a. To determine the number and distribution of large animals
- b. To map out various human activities to assess current pressures on wildlife conservation
- c. To document the number and distribution of livestock in relation to large mammals in the ecosystem
- d. To establish spatial distribution of threats to wildlife due to anthropogenic activities
- e. To interpret the information obtained and deduce sound management decisions to guide management of wildlife in the ecosystem
- f. Provide baseline wildlife data for the National Wildlife census





Figure 40: Map of the study area showing the extent, county boundaries, and the KWS conservations regions.





CHAPTER

4

Conclusions and Recommendations

4.1 | CONCLUSIONS

The national elephant population has increased from 32,214 animals in 2014 to the current population of 36,169 animals in 2021. This represents a 12% population increase in 7 years or 1.75% annual increase over the period. The 2014 period marked the peak of poaching in Kenya. During the period, elephant population was adversely affected by the poaching. The population reduced from 35,588 elephants in 2012 to 32,214 elephants in 2014. Since then (2014), the elephant population has increased. Therefore, efforts put in place by the Government to curb elephant poaching in different ecosystems continue to be productive. These efforts should be continued to further sustain future elephant population growth and range expansion.

The national giraffe population also recorded growth from about 23,000 animals in 2019 to about 34,240 animals in 2021, which represents about 49% increase in three years. However, this increase is attributed to inclusion of more updated data of reticulated giraffe in northern Kenya (Wajir, Mandera, Garissa and Turkana Counties) in 2021 as the data used in 2019 for these counties was for 2011. However, it is important to note that there has been a notable increase of all the three species of giraffe found in Kenya. Efforts to increase penalties on crimes related to threatened species appear to be bearing fruits. Such efforts should be sustained for long term conservation and management of this species.

The three key ecosystems for buffalo include Maasai Mara ecosystem, Tsavo ecosystem, Lake Nakuru National Park and Laikipia-Samburu-Marsabit ecosystem. The Maasai Mara ecosystem recorded the highest population of buffalo in the country (n=11, 604 buffaloes), which represents about 28% of the total buffalo population in Kenya. This is followed by Tsavo ecosystem (19%), Lake Nakuru National Park (15%) and then Laikipia-Samburu-Marsabit ecosystem (15%). These four ecosystems account for about 78% of the total Kenya's buffalo population. Efforts to sustain these populations through veterinary interventions to prevent disease outbreaks, climate mitigation measures to evade drought impacts on their population, enhance security surveillance to prevent poaching for bush meat and use of conservancy concept outside protected areas are necessary.

Our results revealed that Kenya has a population of 473 sitatunga out of which 424 and 49 individuals were recorded in Western and Central Rift Conservation Areas respectively. It is important to note that these populations occur in unprotected wetlands across their range. It is only in Saiwa Swamp National Park and Kitale Nature Conservancy where this species occur in a protected areas. It is therefore important to engage with communities and County Governments where these species occur to explore options of making them acquire some conservation status. Community conservancies and County Reserves can be established in such range sites by communities and County Governments.

Special attention should also be given to species with less than 100 individuals (e.g., roan and sable antelope and mountain bongo) so that their conservation status is enhanced. Interventions like developing of predator free sanctuaries and intensive management need to be explored to save these species from local extinction. A special example that has proved that such species populations can grow is the hirola in Ishaqbin Sanctuary whose hirola population has continued to grow since its establishment. This has improved the National Hirola population which stands at 497 animals in 2021. In addition, options to enhance the population is to import individuals from other wild and captive source sites outside Kenya. This will promote revival of the species and genetic mixing to improve survival of new progeny.

There has been an influx of livestock into the key wildlife ecosystem like Laikipia-Samburu-Meru-Marsabit, Tsavo, Maasai Mara and Lamu-Lower Garissa. This scenario will possibly affect the wildlife species negatively as their habitats become encroached and competition for resources (water, space and forage) increase. As such, displaced of wildlife is likely to occur as they avoid competition with the livestock. This was observed in Laikipia-Samburu-Marsabit-Meru Ecosystem, where it is believed elephants relocated to the hilly areas in the ecosystem, which made it difficult for the census team to sight and count them leading to an overall recording of less population than was recorded in 2017. Such incursions also fuel poaching as most herdsmen are armed with automatic weapons.

Also, there is continued pressure of human activities on wildlife and its habitats within most of the ecosystems. It is unlikely that majority of the wildlife species would co-exist with livestock or in areas with high density of human settlements. Conflicts like crop raids, human-deaths and injury, property destruction and disruption of community social and economic life are destined to ensue in new crop farming and settlement areas. Efforts to develop spatial plans by county governments will help separate humans and wildlife or promote human-wildlife co-existence programs thereof.

Lastly, aerial and ground census methodologies are expensive and labour intensive to implement at national scale. Alternative new technologies that utilize artificial intelligence, operating in a more intensely targeted way

over a much smaller landscapes, have recently been tried to count wildlife. For example application of remote sensing combined with artificial intelligence software is promising as the future of wildlife censuses. It is likely, given their agreement with historic estimates that these will be more suited to the long term monitoring of wildlife in future. However, until newer methods can deliver a calibrated trend for population status monitoring and management uses, the current aerial and ground census methodologies are still satisfactory for these purposes.

4.1 | RECOMMENDATIONS

From our findings we recommend the following:

1. The National Wildlife Census to be conducted after every three years in line with wildlife ecological cycles. In this regard, a budgetary allocation by the National Treasury during the budget cycle every three years is critical.
2. An annual budget is required to support annual and active monitoring of rare and endemic species such as roan antelope, sable antelope, mountain bongo, rhino, Grevy's zebra, hirola and sitatunga
3. An immediate budget support is required to establish status of species currently threatened by illegal trade and bush meat such as pangolins, dik diks and gazelles and those species that were not covered during this census such as leopard, small carnivores and non-human primates
4. There is need for review of legislation to recognize community conservancies as protected areas as they constitute important wildlife range
5. To improve the ecological integrity of dormant parks and reserves, there is need to enhance public-private partnerships
6. A budgetary support is required to establish a data portal and access mechanism at the Wildlife Research and Training Institute to inform wildlife management as part of its mandate
7. For the endangered, rare and endemic species such as the black rhino, mountain bongo, roan antelope, sable antelope among others, there is need for budgetary allocation to support development and implementation of their recovery plans.
8. Deliberate efforts to secure existing rhino population and additional space for rhinos due to the huge investment costs, security and management requirements
9. Invest in new innovations, modern census equipment and software to improve efficiency in undertaking wildlife censuses.
10. The County Governments to incorporate the census findings in the development of the County Integrated Development Plans (CIDP) and spatial plans.
11. The WRTI to pilot models of wildlife utilization programme as part of management of common species to enhance benefits to land owners who live with wildlife.
12. Fast track the full operationalisation of the WRTI which is the corporate body mandated to undertake and coordinate wildlife research.
13. Undertake a national classification of species with low populations in consultation with IUCN.

REFERENCES

African Elephant database. Occasional paper Series of the IUCN Species Survival Commission, No. 60 IUCN/SSC African Elephant Specialist Group, IUCN, Glad, Switzerland.

Barnes, R.F.W., Beardsley, K., Michelmores, F., Barnes, K.L., Alers, M.P.T. & Blom, A. (1997) Estimating forest elephant numbers with dung counts and a geographic information system. *J. Wildl. Manage.* 61, 1384–1393.

Chase, M., Schlossberg, S., Kenana, L., Mukeka, J., Kanga, E., Kiambi, S., Ngene, S. and Omondi, P. (2015) Current status of key savanna elephant populations in Kenya. A Technical Report for Kenya Service, Nairobi, Kenya.

Douglas-Hamilton I. (1996) Counting elephants from the air - total counts. In: *Studying Elephants*. Ed. K. Kangwana. AWF Technical Handbook series, African Wildlife Foundation, Nairobi, Kenya.

GoK (2010). Government of Kenya, Tourism Act 2010. Nairobi: Government Printer.

KWS, TAWIRI & AWF (2015) Cross border aerial census manual, 1st edition 2016. KWS Service and TAWIRI, Nairobi & Arusha, Kenya & Tanzania.

Kibara ON, Odhiambo NM, Njuguna JM (2012). Tourism and economic growth in Kenya: An empirical investigation. *The International Business Economics Research Journal (Online)*, 11(5):517. Available at: <https://doi.org/10.19030/iber.v11i5.6970>

Litoroh, M., Ihwagi, F.W., Mayiinda, R., Bernard, J. and Douglas-Hamilton, I. (2010) Total Aerial Count of Elephants in Laikipia-Samburu Ecosystem in November 2008. Technical Report for Kenya Wildlife Service, Nairobi, Kenya.

Ngene, S., Mukeka, J., Ihwagi, I., Mathenge, J., Wandera, A., Anyona, G., Tobias, N., Kawira, L., Muthuku, I., Kathiwa, J., Gacheru, P., Davidson, Z., King, J. and Omondi, P. (2013) Total aerial count of elephants, Grevy's zebra and other large mammals in Laikipia-Samburu-Marsabit Ecosystem in (November 2012). A Technical Report for Kenya Wildlife Service, Nairobi, Kenya.

Mwangi, P., Ngene, S. and Esau, K. (2007) Wet Season Aerial Count of Large Mammals in the Meru Conservation Area (MCA). A Technical Report for Kenya Wildlife Service, Nairobi, Kenya.

Omondi, P., Bitok, E., Kahindi, O. and Mayiinda R. (2002). Total Aerial Count of Elephants in Laikipia-Samburu Ecosystem. Kenya Wildlife Service Report, Nairobi, Kenya.

Thouless, C., Dublin, H.T., Blanc, J.J., Skinner, D.P., Daniel, T.E., Taylor, R.D., Maisels, F., Fredrick, H.L. and Bouche, P. (2016) African Elephant Status Report 2016: an update from the
Vanleeuwe, H. (2008) Counting elephants in Montane forests: some sources of error. *Afr. J. Ecol.*, 47, 164–174.

Thouless, C., King, J., Kahumbu, P. and Douglas-Hamilton, I. (2008). The Status of Kenya's Elephants 1990 to 2002. Kenya Wildlife Service and Save the Elephants Joint Report, Kenya.

Zar, J.H. (1996) Biostatistical Analysis. Prentice Hall, New Jersey.









REPUBLIC OF KENYA

Ministry of Tourism & Wildlife

CONTACT US

KENYA WILDLIFE SERVICE



KWS Headquarters
Langata Road



+254 (20) 2379407 | +254 (20) 2379408
| +254 (0) 726 610508/9
Toll free: 0800597000



customerservice@kws.go.ke
www.kws.go.ke

[0800 978 9914 40 397 8]



9 789914 403978



REPUBLIC OF KENYA

Wildlife Research & Training Institute



Wildlife Research & Training Institute (WRTI)
P.O. Box 842-20117, Naivasha



+254 700 000 321



director@wrti.go.ke
wrti.kenya@gmail.com